

Multiresolution Analysis for Compactly Supported Interpolating Tensor Product Wavelets

Tommi Höynälänmaa
Institute of Physics, Tampere University of Technology,
P. O. Box 692, FI-33101 Tampere, Finland

We construct multidimensional interpolating tensor product multiresolution analyses (MRA) of the function spaces $C_0(\mathbb{R}^n, K)$, $K = \mathbb{R}$ or $K = \mathbb{C}$, consisting of real or complex valued functions on \mathbb{R}^n vanishing at infinity and the function spaces $C_u(\mathbb{R}^n, K)$ consisting of bounded and uniformly continuous functions on \mathbb{R}^n . We also construct an interpolating dual MRA for both of these spaces. The theory of the tensor products of Banach spaces is used. We generalize the Besov space norm equivalence from the one-dimensional case to our n -dimensional construction.

Keywords: interpolating wavelets, multivariate wavelets, multiresolution analysis, tensor product, injective tensor norm, projective tensor norm, Besov space

MSC: 46A32, 46B28, 15A69, 46E10

1. Introduction

Chui and Li [6] have constructed a one-dimensional MRA of function space $C_u(\mathbb{R})$ (bounded and uniformly continuous complex valued functions on \mathbb{R}) for interpolating wavelets. Donoho [17] has derived convergence results for interpolating wavelets on space $C_0(\mathbb{R})$. Goedecker's book [20] contains also some material about multidimensional interpolating wavelets. Goedecker [20] uses the term interpolating wavelets to mean Deslauriers-Dubuc wavelets whereas some other authors such as Chui and Li [6] use the term to mean roughly wavelet families whose mother scaling function has the cardinal interpolation property $\varphi(k) = \delta_{k,0}$ for all $k \in \mathbb{Z}$ (one-dimensional case). We follow the latter convention in this article. Chui and Li [5] have also constructed a general framework for multivariate wavelets. However, the theory in that article uses function space $L^2(\mathbb{R}^n)$ and so it is unsuitable for the approach in this article. Dubuc and Deslauriers [12, 18] have investigated interpolation processes related to the Deslauriers-Dubuc fundamental functions. Han and Jia [23] have discussed fundamental functions (see the article for a definition) satisfying the cardinal interpolation property. Numerical values for the wavelet filters of the Deslauriers-Dubuc wavelets have been given in [20] and [30].

Theory for orthonormal wavelets has been developed e.g. by Daubechies [8, 9], Meyer [38], and Wojtaszczyk [45]. Kovačević and Sweldens [30] have investigated the use of digital filters for multidimensional biorthogonal wavelets. Lewis [36] has constructed a MRA of function space $L^2(\mathbb{R})$ for interpolating wavelets in one dimension. Lawton et al. [35] have presented conditions for the refinement mask of an orthonormal MRA. Ji and Shen [27] have given a condition for the refinement

mask to be interpolatory and a condition for a dual refinement mask. Krivoshein and Skopina [31] have presented convergence results for frame-like wavelets that are not frames. Karakaz'yan et al. [29] have described symmetric interpolatory masks generating dual compactly supported wavelet systems and they have also given formulas for dual refinement masks. Shui et al. [43] have shown how to construct M -band wavelets with all the following properties: compact support, orthogonality, linear-phase, regularity, and interpolation. This kind of scaling functions exist only when $M \geq 4$. Dahlke et al. [7] have developed a new method to construct higher dimensional scaling functions. Their method is based on solutions to specific Lagrange interpolation problems by polynomials. Han [22] has investigated symmetries of refinable functions. DeVore et al. [13] have developed some theory for multidimensional orthonormal MRA of space $L^p(\mathbb{R}^d)$, $0 < p < \infty$, $d \in \mathbb{Z}_+$. He and Lai [25] have constructed nonseparable box spline wavelets on Sobolev spaces $H^s(\mathbb{R}^2)$. Wavelets have also been discussed in [24].

The theory about tensor products of Banach spaces has been presented e.g. in [41]. We use the notation from [41] for Banach space tensor products in this article. Schaefer's book [42] contains some material about tensor products of topological vector spaces. Book [37] contains also an introduction to tensor products of Banach spaces. Domański et al. [15, 16] have done some research on them. Michor [39] has represented tensor products of Banach spaces using category theory. Grothendieck [21] gives a classical presentation for tensor products of locally convex spaces. Theory for tensor products of Banach spaces can also be found in book [11]. Kustermans and Vaes [32] have developed some theory for space $C_0(G)$ where G is a compact or a locally compact group and for the tensor products of $C_0(G)$. Daws [10] has presented some material on tensor products of Banach algebras. Glöckner [19] has shown that the tensor products of topological vector spaces are not associative. Reinov [40] has investigated Banach spaces without the approximation property. Brodzki and Niblo [4] have done some research on the rapid decay property of discrete groups and the metric approximation property.

An introduction to Besov spaces can be found e.g. in [3]. An introduction to Besov spaces using the Fourier transform based definition of these spaces can be found in [44]. DeVore and Popov [14] have investigated the connection of Besov spaces with the dyadic spline approximation and interpolation of Besov spaces. Kyriazis and Petrushev [33] have given a method for the construction of unconditional bases for Triebel-Lizorkin and Besov spaces. The relationship between orthonormal wavelets and Besov spaces has been discussed by Meyer [38]. Almeida [1] has investigated wavelet bases in (generalized) Besov spaces.

Section 2 introduces some definitions used in the rest of this article. We give some general definitions needed by the MRA's in section 3. A multivariate MRA of $C_u(\mathbb{R}^n, K)$, $K = \mathbb{R}$ or $K = \mathbb{C}$, is constructed in section 4. A multivariate MRA of $C_0(\mathbb{R}^n, K)$, $K = \mathbb{R}$ or $K = \mathbb{C}$, is constructed in section 5. The interpolating dual MRA is presented in section 6. The Besov space norm equivalence from Donoho [17] is generalized for the n -dimensional interpolating MRA's in section 7. A longer

version of this article with more detailed proofs can be found at [26].

2. Preliminaries

2.1. General

The set of all positive real numbers is denoted by \mathbb{R}_+ and the extended real line by \mathbb{R}_* . The set of all positive integer numbers is denoted by \mathbb{Z}_+ . We define $\mathbb{R}_0 := \{x \in \mathbb{R} : x \geq 0\}$ and $\mathbb{R}_{0*} := \{x \in \mathbb{R}_* : x \geq 0\}$. When A and B are arbitrary sets, f is a function from A into B , and $X \subset A$ the image of X under f is denoted by $f[X]$. The set-theoretic support of a function $f : X \rightarrow \mathbb{C}$ where X is a set is denoted by $\text{supp}_{\text{set}} f$. The topological support of a function $f : T \rightarrow \mathbb{C}$ where T is a topological space is denoted by $\text{supp} f$. When E is a metric space, $x \in E$, and $r \in \mathbb{R}_+$ the closed ball of radius r centred at x is denoted by $\overline{B}_E(x; r)$.

Definition 2.1. When $n \in \mathbb{Z}_+$ define

$$Z(n) := \{k \in \mathbb{Z}_+ : k \leq n\}.$$

We define the differences $\Delta_{\mathbf{h}}^m$ and $\Delta_{\mathbf{h}}$ as in [44] and [2, def. V.4.1].

Definition 2.2. Define

$$(\Delta_{\mathbf{h}}^1 f)(\mathbf{x}) := (\Delta_{\mathbf{h}} f)(\mathbf{x}) := f(\mathbf{x} + \mathbf{h}) - f(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{h} \in \mathbb{R}^n$, and $f \in \mathbb{C}^{\mathbb{R}^n}$ and

$$\Delta_{\mathbf{h}}^m f := \Delta_{\mathbf{h}}(\Delta_{\mathbf{h}}^{m-1} f)$$

for all $m \in \mathbb{N} + 2$, $\mathbf{h} \in \mathbb{R}^n$, and $f \in \mathbb{C}^{\mathbb{R}^n}$.

Definition 2.3. Let $K = \mathbb{R}$ or $K = \mathbb{C}$ and $n \in \mathbb{Z}_+$. Let f be a function from \mathbb{R}^n into K . Define

$$N_{\text{cover}}(f) := \max_{\mathbf{x} \in \mathbb{R}^n} \#\{\mathbf{k} \in \mathbb{Z}^n : f(\mathbf{x} - \mathbf{k}) \neq 0\}.$$

Definition 2.4. Let E and F be normed vector spaces. When f is a compactly supported function from E into F define

$$r_{\text{supp}}(f) := \inf\{r \in \mathbb{R}_0 : \text{supp}_{\text{set}} f \subset \overline{B}_E(0; r)\}.$$

Definition 2.5. Let $n \in \mathbb{Z}_+$. Define

$$I_{\text{trans}}(f, \mathbf{x}) := \{\mathbf{k} \in \mathbb{Z}^n : f(\mathbf{x} - \mathbf{k}) \neq 0\}$$

for all $f \in \mathbb{C}^{\mathbb{R}^n}$ and $\mathbf{x} \in \mathbb{R}^n$.

2.2. Sequences and Cartesian Products

Definition 2.6. When $n \in \mathbb{Z}_+$ define

$$\mathbf{0}_n := (0)_{k \in Z(n)}.$$

Definition 2.7. When $n \in \mathbb{Z}_+$ define $J_+(n) := \{0, 1\}^n \setminus \{\mathbf{0}_n\}$.

Definition 2.8. Define $\check{e}_k := (\delta_{j,k})_{j \in \mathbb{Z}}$ for all $k \in \mathbb{Z}$. When $n \in \mathbb{Z}_+$ define $\mathbf{e}_k^{[n]} := \mathbf{e}_k^{Z(n)}$ for all $k \in Z(n)$.

2.3. Vector Spaces

Suppose that A and B are topological vector spaces. We define $\mathcal{L}(A, B)$ to be the set of all continuous linear functions from A into B .

The term *operator* shall mean a bounded linear function from a normed vector space into another one. The term *projection* shall mean a linear projection of a vector space onto a vector space. The topological dual of a Banach space A is denoted by A^* . Unless otherwise stated A^* is equipped with the norm topology.

When A and B are Banach spaces we use the notation $A \subset_1 B$ to mean that A is isometrically embedded in B , i.e. $A \subset B$ and the inclusion map is distance preserving. When A and B are topological vector spaces we define $A =_{\text{tvs}} B$ to be true iff A and B are equal topological vector spaces. When B is a Banach space we use the notation $A \subset_{\text{c.s.}} B$ to mean that A is a closed subspace of B . When E is a Banach space, $A \subset E$, and $B \subset E^*$ we define $B \perp A$ to be true iff $\langle \tilde{b}, a \rangle = 0$ for all $a \in A$ and $\tilde{b} \in B$.

When E is a normed vector space and $x \in E$ we may use the notation $\|x|E\|$ to mean the norm of x in E . When B is a normed vector space and we write $A :=_{\text{n.s.}} \{x \in B : P(x)\}$ we assume that $\|x|A\| := \|x|B\|$ for each $x \in A$.

When E is a Banach space and A and B open or closed subspaces of E the topological direct sum of A and B is denoted by $A \dot{+} B$.

Definition 2.9. Let E be a set and A be a vector space so that $A \subset E$. Suppose that $\|\cdot\|_A : E \rightarrow \mathbb{R}_{0*}$ is a norm on A . We say that the norm $\|\cdot\|_A$ characterizes A on E iff $\|x\|_A < \infty$ for all $x \in A$ and $\|y\|_A = \infty$ for all $y \in E \setminus A$.

2.4. Function Spaces and Sequence Spaces

When $p \in [1, \infty]$, B is a Banach space, and I a denumerable set we denote the l^p space consisting of a subset of B^I by $l^p(I, B)$. As usual, $l^p(I) :=_{\text{n.s.}} l^p(I, \mathbb{C})$. When $K = \mathbb{R}$ or $K = \mathbb{C}$, $n \in \mathbb{Z}_+$, and $p \in [1, \infty]$ we denote the L^p space consisting of a subset of the Borel functions from \mathbb{R}^n into K by $L^p(\mathbb{R}^n, K)$. As usual, $L^p(\mathbb{R}^n) :=_{\text{n.s.}} L^p(\mathbb{R}^n, \mathbb{C})$. When X and Y are Borel spaces define $\text{bor}(X, Y)$ to be the set of all Borel functions from X into Y .

Let $K = \mathbb{R}$ or $K = \mathbb{C}$. When T is a topological space we denote the space of K -valued bounded and continuous functions on T by $C_b(T, K)$. When E is a metric

space the space of K -valued bounded and uniformly continuous functions on E is denoted by $C_u(E, K)$. When T is a locally compact Hausdorff space the space of K -valued bounded and continuous functions vanishing at infinity [17, 34] is denoted by $C_0(T, K)$. The space of K -valued compactly supported and continuous functions on T is denoted by $C_{\text{com}}(T, K)$.

Definition 2.10. Let $K = \mathbb{R}$ or $K = \mathbb{C}$ and I be a denumerable set. Define

$$c_0(I, K) := \{(a_\lambda)_{\lambda \in I} : (a_{\sigma(k)})_{k=0}^\infty \in c_0(\mathbb{N}, K) \text{ for some bijection } \sigma : \mathbb{N} \rightarrow I\}$$

$$\|\mathbf{a}\|_{c_0(I, K)} := \|\mathbf{a}\|_\infty, \quad \mathbf{a} \in c_0(I, K).$$

Banach space $c_0(I, K)$ is isometrically isomorphic to $c_0(\mathbb{N}, K)$.

Definition 2.11. Let $K = \mathbb{R}$ or $K = \mathbb{C}$ and $n \in \mathbb{Z}_+$. Let E be a closed subspace of $C_b(\mathbb{R}^n, K)$ for which the following condition is true:

$$\forall f \in C_b(\mathbb{R}^n, K), c \in \mathbb{R}_+, \mathbf{d} \in \mathbb{R}^n : f \in E \iff f(c \cdot -\mathbf{d}) \in E.$$

Let $a \in \mathbb{R}_+$, $\mathbf{b} \in \mathbb{R}^n$, and $\tilde{f} \in E^*$. The a -dilatation and \mathbf{b} -translation of \tilde{f} , denoted by $\tilde{f}(a \cdot -\mathbf{b})$, is defined by [6]

$$\langle \tilde{f}(a \cdot -\mathbf{b}), f \rangle := \frac{1}{a^n} \left\langle \tilde{f}, f \left(\frac{\cdot + \mathbf{b}}{a} \right) \right\rangle$$

for all $f \in E$.

Definition 2.12. Let $f \in \mathbb{C}^{\mathbb{R}^n}$, $r_1 \in \mathbb{R}_+$, and $\tilde{f} \in C(\overline{B_{\mathbb{R}^n}(0; r_1)})^*$. Let $m \in \mathbb{N}$. We say that pair (\tilde{f}, f) spans all the polynomials of degree at most m iff

$$\sum_{\mathbf{k} \in \mathbb{Z}^n} \left\langle \tilde{f}, p(\cdot + \mathbf{k})|_{\overline{B_{\mathbb{R}^n}(0; r_1)}} \right\rangle f(\mathbf{x} - \mathbf{k}) = p(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{R}^n$ and for all polynomials p of n variables that are of degree at most m .

We define the modulus of continuity in the standard way [2, def. V.4.2]:

Definition 2.13. When $n \in \mathbb{Z}_+$, $m \in \mathbb{Z}_+$, and $p \in [1, \infty]$ define

$$\omega_p^m(f; t) := \sup\{\|\Delta_{\mathbf{h}}^m f\|_{L^p(\mathbb{R}^n)} : \mathbf{h} \in \overline{B_{\mathbb{R}^n}(0; t)}\}$$

and

$$\omega(f; t) := \omega_\infty^1(f; t)$$

for all $f \in \text{bor}(\mathbb{R}^n, \mathbb{C})$ and $t \in \mathbb{R}_0$.

The Besov spaces on \mathbb{R}^n are denoted by $B_{p,q}^s(\mathbb{R}^n)$. We have $\mathcal{Z}^s(\mathbb{R}^n) =_{\text{tvs}} B_{\infty, \infty}^s(\mathbb{R}^n)$ for all $s \in \mathbb{R}_+$ where $\mathcal{Z}^s(\mathbb{R}^n)$ are the Hölder-Zygmund spaces [44]. We also have $C^s(\mathbb{R}^n) =_{\text{tvs}} \mathcal{Z}^s(\mathbb{R}^n) =_{\text{tvs}} B_{\infty, \infty}^s(\mathbb{R}^n)$ when $s \in \mathbb{R}_+ \setminus \mathbb{Z}_+$ where $C^s(\mathbb{R}^n)$, $s \in \mathbb{R}_+ \setminus \mathbb{Z}_+$, are the Hölder spaces on \mathbb{R}^n [44]. When $m \in \mathbb{N}$ we denote the Banach space of functions with bounded and uniformly continuous partial derivatives up to m th degree equipped with the usual norm by $C^m(\mathbb{R}^n)$, see [44].

Definition 2.14. Let $n \in \mathbb{R}_+$, $s \in \mathbb{R}_+$, $p \in [1, \infty]$, and $q \in [1, \infty]$. Let $m \in \mathbb{Z}_+$ and $m > s$. Define

$$\|f\|_{B_{p,q}^s(\mathbb{R}^n);m}^{(c)} := \|f\|_{L^p(\mathbb{R}^n)} + \left\| t \in]0, 1[\mapsto t^{-s-\frac{1}{q}} \omega_p^m(f; t) \right\|_{L^q(]0,1])}$$

for all $f \in \text{bor}(\mathbb{R}^n, \mathbb{C})$.

Norm $\|\cdot\|_{B_{p,q}^s(\mathbb{R}^n);m}^{(c)}$ is an equivalent norm on $B_{p,q}^s(\mathbb{R}^n)$ for the range of parameters given in Definition 2.14. Norm $\|\cdot\|_{B_{p,q}^s(\mathbb{R}^n);m}^{(c)}$ characterizes the Besov space $B_{p,q}^s(\mathbb{R}^n)$ on $\text{bor}(\mathbb{R}^n, \mathbb{C})$ for the given range of parameters. Note that $L^p(\mathbb{R}^n) \subset S'(\mathbb{R}^n)$ for all $n \in \mathbb{Z}_+$ and $p \in [1, \infty]$. See [2, def. V.4.3].

2.5. Tensor Products

Definition 2.15. When $n \in \mathbb{Z}_+$ define

$$\check{\mathbf{e}}_{\mathbf{k}}^{\otimes} := \bigotimes_{l=1}^n \check{\mathbf{e}}_{\mathbf{k}[l]}.$$

for all $\mathbf{k} \in \mathbb{Z}^n$.

When $n \in \mathbb{Z}_+$, α is a uniform crossnorm, and A_1, \dots, A_n are Banach spaces then the completed tensor product of several Banach spaces is defined by the recursive formula

$$A_1 \hat{\otimes}_{\alpha} \cdots \hat{\otimes}_{\alpha} A_n :=_{\text{n.s.}} (A_1 \hat{\otimes}_{\alpha} \cdots \hat{\otimes}_{\alpha} A_{n-1}) \hat{\otimes}_{\alpha} A_n.$$

The indexed completed tensor product is defined by

$$\bigotimes_{j=1}^n \hat{\otimes}_{\alpha} A_j :=_{\text{n.s.}} \left(\bigotimes_{j=1}^{n-1} \hat{\otimes}_{\alpha} A_j \right) \hat{\otimes}_{\alpha} A_n$$

where $n > 1$. When $n = 1$ define

$$\bigotimes_{j=1}^1 \hat{\otimes}_{\alpha} A_j :=_{\text{n.s.}} A_1.$$

Uniform crossnorms do not generally respect subspaces. However, when X is a closed subspace of a Banach space E , Y is a closed subspace F , and α is a reasonable crossnorm on $E \otimes F$ it is possible to use the norm inherited from $E \otimes_{\alpha} F$ on vector space $X \otimes Y$ [37, chapter 1]. We give the following definition for this kind of construction.

Definition 2.16. Let E and F be Banach spaces and α a norm on $E \otimes F$. Let X be a closed subspace of E and Y a closed subspace of F . Define $X \otimes_{\alpha; E \hat{\otimes}_{\alpha} F} Y$ to be the normed vector space formed by using the norm inherited from $E \hat{\otimes}_{\alpha} F$ in the vector space $X \otimes Y$, i.e. $\|u\|_{X \otimes_{\alpha; E \hat{\otimes}_{\alpha} F} Y} := \alpha_{E,F}(u)$ for all $u \in X \otimes Y$. Define $X \hat{\otimes}_{\alpha; E \hat{\otimes}_{\alpha} F} Y$

to be the closure of $X \otimes_{\alpha; E \hat{\otimes}_{\alpha} F} Y$ in space $E \hat{\otimes}_{\alpha} F$. The notations $X \otimes_{\alpha; E \otimes F} Y$ and $X \hat{\otimes}_{\alpha; E \otimes F} Y$ may also be used for the aforementioned definitions.

Now $X \otimes_{\alpha; E \hat{\otimes}_{\alpha} F} Y$ is a normed subspace of $E \hat{\otimes}_{\alpha} F$ and $X \hat{\otimes}_{\alpha; E \hat{\otimes}_{\alpha} F} Y$ is a closed subspace of $E \hat{\otimes}_{\alpha} F$.

Definition 2.17. Let $n \in \mathbb{Z}_+$. Let A_1, \dots, A_n and B_1, \dots, B_n be Banach spaces so that A_1 is a closed subspace of B_1 and $B_k \otimes A_{k+1}$ is a linear subspace of B_{k+1} for $k \in Z(n-1)$. When $k \geq 2$ define

$$\bigotimes_{(B_j)}^k A_j :=_{\text{n.s.}} \text{clos}_{B_k} T_k$$

where

$$T_k :=_{\text{n.s.}} \left(\bigotimes_{(B_j)}^{k-1} A_j \right) \otimes_{(B_k)} A_k.$$

When $k = 1$ define

$$\bigotimes_{(B_j)}^k A_j :=_{\text{n.s.}} A_1.$$

When the assumptions of Definition 2.17 hold

$$\bigotimes_{(B_j)}^k A_j$$

is a closed subspace of Banach space B_k for all $k \in Z(n)$.

Definition 2.18. Let $n \in \mathbb{Z}_+$. Let A_1, \dots, A_n , B_1, \dots, B_n , E_1, \dots, E_n , and F_1, \dots, F_n be Banach spaces so that

- A_1 is a closed subspace of E_1 and $E_k \otimes A_{k+1}$ is a linear subspace of E_{k+1} for all $k = 1, \dots, n-1$.
- B_1 is a closed subspace of F_1 and $F_k \otimes B_{k+1}$ is a linear subspace of F_{k+1} for all $k = 1, \dots, n-1$.

Suppose that $P_k : A_k \rightarrow B_k$, $k = 1, \dots, n$ are operators. Let $S_1 := P_1$, $T_1 := S_1 = P_1$, and $S_k := T_{k-1} \otimes P_k$ for $k = 2, \dots, n$. When $k \in \{2, \dots, n\}$ and S_k is continuous let

$$T_k : \bigotimes_{(E_l)}^k A_l \rightarrow \bigotimes_{(F_l)}^k B_l,$$

be the unique continuous linear extension of S_k to

$$\bigotimes_{l=1}^k A_l.$$

If $k \in \{2, \dots, n\}$ and S_k is not continuous let $T_k = 0$. When all of the functions S_k , T_k , $k = 1, \dots, n$ are operators define

$$\bigotimes_{k=1}^n P_k = T_n.$$

If any of the functions S_k , T_k , $k = 1, \dots, n$ is not an operator then

$$\bigotimes_{k=1}^n P_k$$

is undefined.

There is an isometric embedding

$$X^* \otimes_{\alpha^s} Y^* \subset_1 (X \otimes_{\alpha} Y)^* \quad (2.1)$$

for tensor norms α and Banach spaces X and Y [41], see also [11]. When $K = \mathbb{R}$ or $K = \mathbb{C}$, $n \in \mathbb{N} + 2$, and X_1, \dots, X_n are closed subspaces of $C_b(\mathbb{R}, K)$ the tensor product $X_1 \hat{\otimes}_{\varepsilon} \dots \hat{\otimes}_{\varepsilon} X_n$ is a closed subspace of $C_b(\mathbb{R}^n, K)$.

3. General Definitions for a Compactly Supported Interpolating MRA

3.1. Mother Scaling Function

Definition 3.1. Let $K = \mathbb{R}$ or $K = \mathbb{C}$ and $n \in \mathbb{Z}_+$. A compactly supported interpolating mother scaling function is a function $\varphi \in C_{\text{com}}(\mathbb{R}^n, K)$ satisfying the following conditions:

(MSF.1)

$$\forall \mathbf{k} \in \mathbb{Z}^n : \varphi(\mathbf{k}) = \delta_{\mathbf{k},0}$$

(MSF.2)

$$\forall \mathbf{x} \in \mathbb{R}^n : \varphi(\mathbf{x}) = \sum_{\mathbf{k} \in \mathbb{Z}^n} \varphi\left(\frac{\mathbf{k}}{2}\right) \varphi(2\mathbf{x} - \mathbf{k}).$$

The Deslauriers-Dubuc fundamental functions satisfy these conditions [17].

3.2. General Definitions for the Univariate MRA's

We shall denote the function space for which the MRA is defined by E in this section. We have either $E = C_u(\mathbb{R}, K)$ or $E = C_0(\mathbb{R}, K)$ where $K = \mathbb{R}$ or $K = \mathbb{C}$. We shall assume that $\varphi \in C_{\text{com}}(\mathbb{R}, K)$ is a compactly supported interpolating mother scaling function throughout this subsection.

Definition 3.2. Define function $\psi \in C_{\text{com}}(\mathbb{R}, K)$ by

$$\psi(x) := \varphi(2x - 1)$$

for all $x \in \mathbb{R}$. Function ψ is called the mother wavelet.

Definition 3.3. Define

$$\varphi_{j,k} := \varphi(2^j \cdot -k)$$

and

$$\psi_{j,k} := \psi(2^j \cdot -k)$$

for all $j \in \mathbb{Z}$ and $k \in \mathbb{Z}$.

Definition 3.4. Define

$$\zeta_s := \begin{cases} \varphi; & s = 0 \\ \psi; & s = 1 \end{cases}$$

where $s \in \{0, 1\}$.

Definition 3.5. When $k \in \mathbb{Z}$ define

$$\begin{aligned} h_k &:= \varphi\left(\frac{k}{2}\right) \\ g_k &:= \delta_{k,1} \\ \tilde{h}_k &:= \delta_{k,0} \\ \tilde{g}_k &:= (-1)^{k-1} h_{1-k}. \end{aligned}$$

Definition 3.6. Define $\tilde{\varphi} := \delta \in E^*$ and $\tilde{\psi} \in E^*$ by

$$\tilde{\psi} := 2 \sum_{k \in \mathbb{Z}} \tilde{g}_k \tilde{\varphi}(2 \cdot -k). \quad (3.1)$$

Define $\tilde{\varphi}_{j,k} := 2^j \tilde{\varphi}(2^j \cdot -k)$ and $\tilde{\psi}_{j,k} := 2^j \tilde{\psi}(2^j \cdot -k)$ where $j, k \in \mathbb{Z}$.

As φ is compactly supported only a finite number of numbers h_k , $k \in \mathbb{Z}$, and the other three filters defined by Definition 3.5 are nonzero. Consequently the series in Equation (3.1) has a finite number of nonzero terms.

Definition 3.7. Define

$$\tilde{\zeta}_s := \begin{cases} \tilde{\varphi}; & s = 0 \\ \tilde{\psi}; & s = 1 \end{cases}$$

where $s \in \{0, 1\}$.

3.3. General Definitions for Multivariate MRA's

We will assume that $n \in \mathbb{Z}_+$ and either $K = \mathbb{R}$ or $K = \mathbb{C}$ throughout this subsection. We will also assume that $E =_{\text{n.s.}} C_u(\mathbb{R}, K)$ or $E =_{\text{n.s.}} C_0(\mathbb{R}, K)$. Furthermore, $\varphi \in C_{\text{com}}(\mathbb{R}, K)$ shall be a compactly supported interpolating mother scaling function throughout this subsection.

We set

$$F :=_{\text{n.s.}} \begin{cases} C_u(\mathbb{R}^n, K); & E = C_u(\mathbb{R}, K) \\ C_0(\mathbb{R}^n, K); & E = C_0(\mathbb{R}, K) \end{cases}$$

Definition 3.8. Define function $\varphi^{[n]} \in C_{\text{com}}(\mathbb{R}^n, K)$ by

$$\varphi^{[n]} := \bigotimes_{k=1}^n \varphi.$$

Function $\varphi^{[n]}$ is called an n -dimensional tensor product mother scaling function generated by φ . Define also $\varphi_{j,\mathbf{k}}^{[n]} := \varphi^{[n]}(2^j \cdot -\mathbf{k})$ where $j \in \mathbb{Z}$ and $\mathbf{k} \in \mathbb{Z}^n$.

Function $\varphi^{[n]}$ is a compactly supported interpolating mother scaling function on \mathbb{R}^n .

Definition 3.9. When $\mathbf{s} \in \{0, 1\}^n$ define function $\psi_{\mathbf{s}}^{[n]} \in C_{\text{com}}(\mathbb{R}^n, K)$ by

$$\psi_{\mathbf{s}}^{[n]} := \bigotimes_{k=1}^n \zeta_{\mathbf{s}[k]}$$

and $\psi_{\mathbf{s},j,\mathbf{k}}^{[n]} := \psi_{\mathbf{s}}^{[n]}(2^j \cdot -\mathbf{k})$.

The domain of the Dirac δ functional varies in this article. I.e. we may keep δ as an element of different dual spaces A^* . When $z_1, \dots, z_m \in \mathbb{R}$ we will identify $\delta(\cdot - z_1) \otimes \dots \otimes \delta(\cdot - z_m)$ with $\delta(\cdot - (z_1, \dots, z_m))$.

Definition 3.10. Define $\tilde{\varphi}^{[n]} \in F^*$ by

$$\tilde{\varphi}^{[n]} := \bigotimes_{l=1}^n \tilde{\varphi}$$

and $\tilde{\varphi}_{j,\mathbf{k}}^{[n]} \in F^*$ by $\tilde{\varphi}_{j,\mathbf{k}}^{[n]} := 2^j \tilde{\varphi}^{[n]}(2^j \cdot -\mathbf{k})$ where $j \in \mathbb{Z}$ and $\mathbf{k} \in \mathbb{Z}^n$. Define also $\tilde{\psi}_{\mathbf{s}}^{[n]} \in F^*$ by

$$\tilde{\psi}_{\mathbf{s}}^{[n]} := \bigotimes_{l=1}^n \tilde{\zeta}_{\mathbf{s}[l]}$$

where $\mathbf{s} \in \{0, 1\}^n$ and $\tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]} \in F^*$ by $\tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]} := 2^j \tilde{\psi}_{\mathbf{s}}^{[n]}(2^j \cdot -\mathbf{k})$ where $\mathbf{s} \in \{0, 1\}^n$, $j \in \mathbb{Z}$, and $\mathbf{k} \in \mathbb{Z}^n$.

Goedecker [20] gives formulas for wavelet filters, too.

Lemma 3.1. *Let $f \in \mathbb{C}$, $m \in \mathbb{N}$, $r_1 \in \mathbb{R}_+$, and $\tilde{f} \in C(\overline{B_{\mathbb{R}}}(0; r_1))^*$. Let*

$$f^{[n]} := \bigotimes_{i=0}^n f$$

and

$$\tilde{f}^{[n]} := \bigotimes_{i=0}^n \tilde{f}.$$

If (\tilde{f}, f) spans all polynomials of degree at most m then $(\tilde{f}^{[n]}, f^{[n]})$ spans all polynomials of degree at most m .

Proof. Let

$$p(\mathbf{x}) := \prod_{i=1}^n p_i(\mathbf{x}[i]) = \prod_{i=1}^n (\mathbf{x}[i])^{\mathbf{a}[i]}$$

where $\mathbf{a}[i] \in Z_0(m)$, $i = 1, \dots, n$. Define $r_2 \in \mathbb{R}_+$ so that $(\overline{B_{\mathbb{R}^n}}(0; r_1))^n \subset \overline{B_{\mathbb{R}^n}}(0; r_2)$. Now

$$\begin{aligned} & \sum_{\mathbf{k} \in \mathbb{Z}^n} \left\langle \tilde{f}^{[n]}(\cdot - \mathbf{k}), p|_{\overline{B_{\mathbb{R}^n}}(\mathbf{k}; r_2)} \right\rangle f^{[n]}(\mathbf{x} - \mathbf{k}) \\ &= \sum_{k_1 \in \mathbb{Z}} \cdots \sum_{k_n \in \mathbb{Z}} \prod_{i=1}^n \left\langle \tilde{f}(\cdot - \mathbf{k}[i]), p_i|_{\overline{B_{\mathbb{R}}}(\mathbf{k}[i]; r_1)} \right\rangle f(\mathbf{x}[i] - \mathbf{k}[i]) \\ &= \prod_{i=1}^n \sum_{k \in \mathbb{Z}} \left\langle \tilde{f}(\cdot - k), p_i|_{\overline{B_{\mathbb{R}}}(k; r_1)} \right\rangle f(\mathbf{x}[i] - k) \\ &= \prod_{i=1}^n p_i(\mathbf{x}[i]) = p(\mathbf{x}) \end{aligned}$$

for all $\mathbf{x} \in \mathbb{R}^n$. □

In particular, if φ spans all polynomials of degree at most m then $\varphi^{[n]}$ spans all polynomials of n variables and of degree at most m .

4. Compactly Supported Interpolating MRA of $C_u(\mathbb{R}^n, K)$

We will assume that $n \in \mathbb{Z}_+$ and $K = \mathbb{R}$ or $K = \mathbb{C}$ throughout this section. We will also assume that $\varphi^{[n]} \in C_{\text{com}}(\mathbb{R}^n, K)$ is an n -dimensional tensor product mother scaling function in this section. Unless otherwise stated, we shall assume that the same values of n , K , and $\varphi^{[n]}$ are used throughout this section. Chui and Li [6] have developed a MRA in the univariate case $C_u(\mathbb{R})$.

Definition 4.1. *Define*

$$\begin{aligned} V_{n,j}^{(u)} &:= \left\{ \mathbf{x} \in \mathbb{R}^n \mapsto \sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{a}[\mathbf{k}] \varphi^{[n]}(2^j \mathbf{x} - \mathbf{k}) : \mathbf{a} \in l^\infty(\mathbb{Z}^n, K) \right\} \quad (4.1) \\ \|f|V_{n,j}^{(u)}\| &:= \|f\|_\infty, \quad f \in V_{n,j}^{(u)} \end{aligned}$$

for all $j \in \mathbb{Z}$.

Definition 4.2. Let spaces $V_{n,j}^{(u)}$, $j \in \mathbb{Z}$, be defined by Definition 4.1. We call $\{V_{n,j}^{(u)} : j \in \mathbb{Z}\}$ an interpolating tensor product MRA of $C_u(\mathbb{R}^n, K)$ generated by $\varphi^{[n]}$ provided that the following conditions are satisfied:

- (MRA1.1) $\forall j \in \mathbb{Z} : V_{n,j}^{(u)} \subset V_{n,j+1}^{(u)}$
 (MRA1.2) $\bigcup_{j \in \mathbb{Z}} V_{n,j}^{(u)} = C_u(\mathbb{R}^n, K)$
 (MRA1.3) $\bigcap_{j \in \mathbb{Z}} V_{n,j}^{(u)} = K$
 (MRA1.4) $\forall j \in \mathbb{Z}, f \in K^{\mathbb{R}^n} : f \in V_{n,j}^{(u)} \iff f(2 \cdot) \in V_{n,j+1}^{(u)}$
 (MRA1.5) $\forall j \in \mathbb{Z}, \mathbf{k} \in \mathbb{Z}^n, f \in K^{\mathbb{R}^n} : f \in V_{n,j}^{(u)} \iff f(\cdot - 2^{-j} \mathbf{k}) \in V_{n,j}^{(u)}$
 (MRA1.6) $\forall \mathbf{k} \in \mathbb{Z}^n : \varphi^{[n]}(\mathbf{k}) = \delta_{\mathbf{k},0}$

Our requirements for the definition of interpolating multiresolution analysis are stricter and simpler than those in [6]. Condition (MRA1.6) is replaced by a weaker condition for φ in [6] but it is possible to construct function φ_L that satisfies condition (MRA1.6) and generates the same subspaces $V_{1,j}^{(u)}$ as function φ .

Under the conditions given at the start of this section ($\varphi^{[n]}$ is an n -dimensional tensor product mother scaling function) the conditions (MRA1.1), (MRA1.4), (MRA1.5), and (MRA1.6) are true. We will show later that conditions (MRA1.2) and (MRA1.3) are true, too.

Spaces $V_{n,j}^{(u)}$, $j \in \mathbb{Z}$, are topologically isomorphic to $l^\infty(\mathbb{Z}^n, K)$. It has been proved in [28] that

$$\sum_{\mathbf{k} \in \mathbb{Z}^n} \varphi^{[n]}(\mathbf{y} - \mathbf{k}) = \widehat{\varphi^{[n]}}(0)$$

for all $\mathbf{y} \in \mathbb{R}^n$. It follows from the cardinal interpolation property (MSF.1) that

$$\sum_{\mathbf{k} \in \mathbb{Z}^n} \varphi^{[n]}(\mathbf{y} - \mathbf{k}) = 1 \quad (4.2)$$

for all $\mathbf{y} \in \mathbb{R}^n$.

Remark 4.1. The series in Equation (4.1) converges (pointwise) for all $\mathbf{a} \in l^\infty(\mathbb{Z}^n, K)$ but the series $\sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{a}[\mathbf{k}] \varphi^{[n]}(2^j \cdot - \mathbf{k})$ need not converge in the norm topology of $C_u(\mathbb{R}^n, K)$. For example, consider sequence $\mathbf{a}[\mathbf{k}] = 1$, $\mathbf{k} \in \mathbb{Z}^n$. Now $\sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{a}[\mathbf{k}] \varphi^{[n]}(2^j \mathbf{x} - \mathbf{k}) = 1$ for all $\mathbf{x} \in \mathbb{R}^n$ but the series $\sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{a}[\mathbf{k}] \varphi^{[n]}(2^j \cdot - \mathbf{k})$ does not converge in the norm topology of $C_u(\mathbb{R}^n, K)$.

Theorem 4.1. We have

$$\bigcap_{j \in \mathbb{Z}} V_{n,j}^{(u)} = K.$$

Proof. The proof is similar to a part of the proof of [6, theorem 3.2]. \square

Definition 4.3. When $j \in \mathbb{Z}$ and $\mathbf{s} \in \{0, 1\}^n$ define

$$W_{n,\mathbf{s},j}^{(u)} := \left\{ \mathbf{x} \in \mathbb{R}^n \mapsto \sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{a}[\mathbf{k}] \psi_{\mathbf{s},j,\mathbf{k}}^{[n]}(\mathbf{x}) : \mathbf{a} \in l^\infty(\mathbb{Z}^n, K) \right\}$$

$$\|f|W_{n,\mathbf{s},j}^{(u)}\| := \|f\|_\infty, \quad f \in W_{n,\mathbf{s},j}^{(u)}.$$

Spaces $W_{n,\mathbf{s},j}^{(u)}$, $\mathbf{s} \in \{0, 1\}^n$, $j \in \mathbb{Z}$, are topologically isomorphic to $l^\infty(\mathbb{Z}^n, K)$.

Definition 4.4. When $\mathbf{s} \in \{0, 1\}^n$ and $j \in \mathbb{Z}$ define

$$\left(Q_{n,\mathbf{s},j}^{(u)} f \right) (\mathbf{x}) := \sum_{\mathbf{k} \in \mathbb{Z}^n} \left\langle \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]}, f \right\rangle \psi_{\mathbf{s},j,\mathbf{k}}^{[n]}(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{R}^n$ and $f \in C_u(\mathbb{R}^n, K)$. When $j \in \mathbb{Z}$ define $P_{n,j}^{(u)} := Q_{n,\mathbf{0},j}^{(u)}$.

Operator $Q_{n,\mathbf{s},j}^{(u)}$ is a continuous projection of $C_u(\mathbb{R}^n, K)$ onto $W_{n,\mathbf{s},j}^{(u)}$ for each $\mathbf{s} \in \{0, 1\}^n$ and $j \in \mathbb{Z}$.

Definition 4.5. When $j \in \mathbb{Z}$ define

$$W_{n,j}^{(u)} :=_{\text{n.s.}} \sum_{\mathbf{s} \in J_+(n)} W_{n,\mathbf{s},j}^{(u)}$$

and

$$Q_{n,j}^{(u)} := \sum_{\mathbf{s} \in J_+(n)} Q_{n,\mathbf{s},j}^{(u)}.$$

Spaces $W_{n,j}^{(u)}$, $j \in \mathbb{Z}$, are topologically isomorphic to $l^\infty(J_+(n) \times \mathbb{Z}^n, K)$. When $j \in \mathbb{Z}$ we have

$$V_{n,j+1}^{(u)} =_{\text{n.s.}} V_{n,j}^{(u)} \dot{+} W_{n,j}^{(u)} =_{\text{n.s.}} V_{n,j}^{(u)} \dot{+} \sum_{\mathbf{s} \in J_+(n)} W_{n,\mathbf{s},j}^{(u)} =_{\text{n.s.}} \sum_{\mathbf{s} \in \{0,1\}^n} W_{n,\mathbf{s},j}^{(u)}. \quad (4.3)$$

Operator $Q_{n,j}^{(u)}$ is a continuous projection of $C_u(\mathbb{R}^n, K)$ onto $W_{n,j}^{(u)}$ for each $j \in \mathbb{Z}$.

We also have $Q_{n,j}^{(u)} = P_{n,j+1}^{(u)} - P_{n,j}^{(u)}$ for all $j \in \mathbb{Z}$.

Theorem 4.2. There exist $c_1 \in \mathbb{R}_+$ and $c_2 \in \mathbb{R}_+$ so that

$$\left\| f - P_{n,j}^{(u)} f \right\|_\infty \leq c_1 \omega(f; 2^{-j} c_2)$$

for all $f \in C_u(\mathbb{R}^n)$ and $j \in \mathbb{Z}$.

Proof. Let $f \in C_u(\mathbb{R}^n)$ and $j \in \mathbb{Z}$. Define $r_1 := r_{\text{supp}}(\varphi^{[n]})$. Let $\mathbf{x} \in \mathbb{R}^n$. Define $I := I_{\text{trans}}(\varphi^{[n]}, 2^j \mathbf{x})$. Now by Equation (4.2) we have

$$\begin{aligned} \left(f - P_{n,j}^{(u)} f \right) (\mathbf{x}) &= \sum_{\mathbf{k} \in \mathbb{Z}^n} \left(f(\mathbf{x}) - f\left(\frac{\mathbf{k}}{2^j}\right) \right) \varphi^{[n]}(2^j \mathbf{x} - \mathbf{k}) \\ &= \sum_{\mathbf{k} \in I} \left(f(\mathbf{x}) - f\left(\frac{\mathbf{k}}{2^j}\right) \right) \varphi^{[n]}(2^j \mathbf{x} - \mathbf{k}) \end{aligned}$$

Let $\ell \in I$. Now $2^j \mathbf{x} - \ell \in \overline{B_{\mathbb{R}^n}}(0; r_1)$ from which it follows that $\mathbf{x} - 2^{-j} \ell \in \overline{B_{\mathbb{R}^n}}(0; 2^{-j} r_1)$. Hence

$$\left| f(\mathbf{x}) - f\left(\frac{\ell}{2^j}\right) \right| \leq \omega(f; 2^{-j} r_1). \quad (4.4)$$

Let $c_1 := N_{\text{cover}}(\varphi^{[n]}) \|\varphi^{[n]}\|_{\infty}$ and $c_2 := r_1 = r_{\text{supp}}(\varphi^{[n]})$. By Equation (4.4) we have

$$\left| \left(f - P_{n,j}^{(u)} f \right) (\mathbf{x}) \right| \leq c_1 \omega(f; 2^{-j} c_2). \quad \square$$

Theorem 4.3. *Let $f \in C_u(\mathbb{R}^n)$. Now*

$$\lim_{j \rightarrow \infty} \left\| f - P_{n,j}^{(u)} f \right\|_{\infty} = 0.$$

Proof. See also [6, theorem 3.2] and [17, theorem 2.4]. Let $j \in \mathbb{Z}$. By Theorem 4.2 we have $\|f - P_{n,j}^{(u)} f\|_{\infty} \leq c_1 \omega(f; 2^{-j} c_2)$ where c_1 and c_2 do not depend on j or f . Since f is uniformly continuous

$$\lim_{t \rightarrow 0} \omega(f; t) = 0$$

and hence

$$\lim_{j \rightarrow \infty} \left\| f - P_{n,j}^{(u)} f \right\|_{\infty} = 0. \quad \square$$

Corollary 4.1.

$$\overline{\bigcup_{j \in \mathbb{Z}} V_{n,j}^{(u)}} = C_u(\mathbb{R}^n, K).$$

Theorem 4.4. *Let $t \in \mathbb{R}_+$. There exists a constant $c_1 \in \mathbb{R}_+$, which may depend on t , so that*

$$\forall f \in C_u(\mathbb{R}^n), j \in \mathbb{Z} : \omega(P_{n,j}^{(u)} f; 2^{-j} t) \leq c_1 \omega(f; 2^{-j} \sqrt{n}).$$

Proof. The proof of case $n = 1$ is similar to the proof in [17, section 7.1]. We will assume that $n \geq 2$ in the sequel. Define bijection $\sigma_c^{[n]} : \mathbb{N} \rightarrow \mathbb{Z}^n$ so that $\|\sigma_c^{[n]}(p+1) - \sigma_c^{[n]}(p)\|_{\infty} = 1$ for all $p \in \mathbb{N}$. It can be proved that this kind of bijection always exists [26]. Define

$$v_{j,l}(\mathbf{x}, \mathbf{h}) := \sum_{p=0}^l \left(\varphi^{[n]}(2^j \mathbf{x} - \sigma_c^{[n]}(p) - \lfloor 2^j \mathbf{x} \rfloor + 2^j \mathbf{h}) - \varphi^{[n]}(2^j \mathbf{x} - \sigma_c^{[n]}(p) - \lfloor 2^j \mathbf{x} \rfloor) \right)$$

where $j, l \in \mathbb{N}$, $\mathbf{x} \in \mathbb{R}^n$, and $\mathbf{h} \in \overline{B_{\mathbb{R}^n}}(0; 2^{-j} t)$. Using Equation (4.2) it can be proved that there exists $c_1 \in \mathbb{R}_+$ so that

$$\sum_{l=0}^{\infty} |v_{j,l}(\mathbf{x}, \mathbf{h})| \leq c_1$$

for all $j \in \mathbb{N}$, $\mathbf{x} \in \mathbb{R}^n$, and $\mathbf{h} \in \overline{B}_{\mathbb{R}^n}(0; 2^{-j}t)$. Let $f \in C_u(\mathbb{R}^n)$, $j_1 \in \mathbb{Z}$, $\mathbf{x}_1 \in \mathbb{R}^n$, and $\mathbf{h}_1 \in \overline{B}_{\mathbb{R}^n}(0; 2^{-j_1}t)$. Let $\mathbf{a} := (f(2^{-j_1}\mathbf{k}))_{\mathbf{k} \in \mathbb{Z}^n}$. Now

$$\begin{aligned} z &:= \left(P_{n,j_1}^{(u)} f \right) (\mathbf{x}_1 + \mathbf{h}_1) - \left(P_{n,j_1}^{(u)} f \right) (\mathbf{x}_1) \\ &= \sum_{l=0}^{\infty} \left(\mathbf{a} \left[\sigma_c^{[n]}(l) \right] - \mathbf{a} \left[\sigma_c^{[n]}(l+1) \right] \right) v_{j_1,l}(\mathbf{x}_1, \mathbf{h}_1) \end{aligned}$$

from which it follows that

$$|z| \leq \left(\sup_{l \in \mathbb{N}} \left| \mathbf{a} \left[\sigma_c^{[n]}(l) \right] - \mathbf{a} \left[\sigma_c^{[n]}(l+1) \right] \right| \right) \cdot \sum_{l=0}^{\infty} |v_{j_1,l}(\mathbf{x}_1, \mathbf{h}_1)| \leq \omega(f; 2^{-j_1}\sqrt{n}) \cdot c_1. \quad \square$$

Theorem 4.5. *Let $j_0 \in \mathbb{Z}$. Suppose that the mother scaling function φ is Lipschitz continuous and let $\varphi^{[n]}$ be the tensor product mother scaling function generated by φ . Then*

$$V_{n,j_0}^{(u)} \dot{+} \sum_{j=j_0}^{\infty} W_{n,j}^{(u)} \neq C_u(\mathbb{R}^n, K).$$

Proof. Let

$$A := V_{n,j_0}^{(u)} \dot{+} \sum_{j=j_0}^{\infty} W_{n,j}^{(u)}.$$

Define function $f \in C_u(\mathbb{R}, K)$ by

$$f(x) := \begin{cases} \sqrt{x}; & x \in [0, 1] \\ -x + 2; & x \in [1, 2] \\ 0; & x \leq 0 \vee x \geq 2 \end{cases}$$

and function $f^{[n]} \in C_u(\mathbb{R}^n, K)$ by

$$f^{[n]} := \bigotimes_{k=1}^n f.$$

Functions f and $f^{[n]}$ are not Lipschitz continuous. As φ is Lipschitz continuous all the functions $P_{n,j}^{(u)}g$ are Lipschitz continuous for each $g \in C_u(\mathbb{R}^n, K)$. A is a locally convex space with an inductive limit topology. It follows from [42, theorem 6.2] that the space A is complete.

Suppose that A would be equal to $C_u(\mathbb{R}^n, K)$ as a set. Then we would have $f^{[n]} \in A$. It follows from the definition of the locally convex direct sum that

$$f^{[n]} \in V_{n,j_0}^{(u)} \dot{+} \sum_{j=j_0}^{j_1} W_{n,j}^{(u)}$$

for some $j_1 \in \mathbb{Z}$, $j_1 \geq j_0$. Now $f^{[n]} = P_{n,j_1+1}^{(u)}f^{[n]}$. It follows that $f^{[n]}$ is Lipschitz continuous, which is a contradiction. Hence A is not equal to $C_u(\mathbb{R}^n, K)$ as a set. \square

The definition of locally convex direct sums from [42] is used here. An example of a Lipschitz continuous mother scaling function is a Deslauriers-Dubuc fundamental function with Hölder regularity greater than 1. By Equation (4.3) and Corollary 4.1 we have

$$C_u(\mathbb{R}^n, K) =_{\text{n.s.}} \text{clos} \left(\bigcup_{l=j_0}^{\infty} \left(V_{n,j_0}^{(u)} + \sum_{j=j_0}^l W_{n,j}^{(u)} \right) \right)$$

for all $j_0 \in \mathbb{Z}$.

5. Compactly Supported Interpolating MRA of $C_0(\mathbb{R}^n, K)$

A MRA of space $C_0(\mathbb{R}^n, K)$ can be developed in the same way as the MRA of space $C_u(\mathbb{R}^n, K)$. Donoho [17] has developed a MRA for the one-dimensional case $C_0(\mathbb{R})$. We will assume that $n \in \mathbb{Z}_+$ and $K = \mathbb{R}$ or $K = \mathbb{C}$ throughout this section. We will also assume that $\varphi^{[n]} \in C_{\text{com}}(\mathbb{R}^n, K)$ is an n -dimensional tensor product mother scaling function in this section. Unless otherwise stated, we shall assume that the same values of n , K , and $\varphi^{[n]}$ are used throughout this section. Note that many of the series that are pointwise convergent with $C_u(\mathbb{R}^n, K)$ converge in the norm in $C_0(\mathbb{R}^n, K)$. We are also able to represent many subspaces and operators related to the MRA of $C_0(\mathbb{R}^n, K)$ as tensor products of the corresponding one-dimensional cases.

Definition 5.1. *Define*

$$V_{n,j}^{(0)} := \left\{ \sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{a}[\mathbf{k}] \varphi^{[n]}(2^j \cdot -\mathbf{k}) : \mathbf{a} \in c_0(\mathbb{Z}^n, K) \right\}$$

$$\|f|V_{n,j}^{(0)}\| := \|f\|_{\infty}, \quad f \in V_{n,j}^{(0)} \quad (5.1)$$

for each $j \in \mathbb{Z}$.

Definition 5.2. *Let spaces $V_{n,j}^{(0)}$, $j \in \mathbb{Z}$, be defined by Definition 5.1. We call $\{V_{n,j}^{(0)} : j \in \mathbb{Z}\}$ an interpolating tensor product MRA of $C_0(\mathbb{R}^n, K)$ generated by $\varphi^{[n]}$ provided that the following conditions are satisfied:*

- (MRA2.1) $\forall j \in \mathbb{Z} : V_{n,j}^{(0)} \subset V_{n,j+1}^{(0)}$
- (MRA2.2) $\bigcup_{j \in \mathbb{Z}} V_{n,j}^{(0)} = C_0(\mathbb{R}^n, K)$
- (MRA2.3) $\bigcap_{j \in \mathbb{Z}} V_{n,j}^{(0)} = \{0\}$
- (MRA2.4) $\forall j \in \mathbb{Z}, f \in K^{\mathbb{R}^n} : f \in V_{n,j}^{(0)} \iff f(2 \cdot) \in V_{n,j+1}^{(0)}$
- (MRA2.5) $\forall j \in \mathbb{Z}, \mathbf{k} \in \mathbb{Z}^n, f \in K^{\mathbb{R}^n} : f \in V_{n,j}^{(0)} \iff f(\cdot - 2^{-j} \mathbf{k}) \in V_{n,j}^{(0)}$
- (MRA2.6) $\forall \mathbf{k} \in \mathbb{Z}^n : \varphi^{[n]}(\mathbf{k}) = \delta_{\mathbf{k},0}$

Note that in this definition the intersection of spaces $V_{n,j}^{(0)}$ is $\{0\}$ instead of \mathbb{C} (all complex valued constant functions on \mathbb{R}^n) as in [6] since here the MRA

is constructed for functions vanishing at infinity. Donoho [17] does not require the mother scaling function φ to be compactly supported but φ has to be of rapid decay in his construction. He also includes requirements for the regularity and polynomial span of φ , which are needed for the norm equivalences to Besov and Triebel-Lizorkin spaces, into the definition of the MRA. Under the conditions given at the start of this section ($\varphi^{[n]}$ is an n -dimensional tensor product mother scaling function) the conditions (MRA2.1), (MRA2.2), (MRA2.3), (MRA2.4), (MRA2.5), and (MRA2.6) are true.

Definition 5.3. *Let $j \in \mathbb{Z}$ and $\mathbf{s} \in \{0, 1\}^n$. Define*

$$W_{n,\mathbf{s},j}^{(0)} := \left\{ \sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{a}[\mathbf{k}] \psi_{\mathbf{s},j,\mathbf{k}}^{[n]} : \mathbf{a} \in c_0(\mathbb{Z}^n, K) \right\}$$

$$\|f\|_{W_{n,\mathbf{s},j}^{(0)}} := \|f\|_{\infty}, \quad f \in W_{n,\mathbf{s},j}^{(0)}.$$

Definition 5.4. *Let $j \in \mathbb{Z}$. Define operator $P_{n,j}^{(0)} : C_0(\mathbb{R}^n, K) \rightarrow V_{n,j}^{(0)}$ by*

$$P_{n,j}^{(0)} f = \sum_{\mathbf{k} \in \mathbb{Z}^n} \langle \tilde{\varphi}_{j,\mathbf{k}}^{[n]}, f \rangle \varphi_{j,\mathbf{k}}^{[n]}$$

for all $f \in C_0(\mathbb{R}^n, K)$. Define operator $Q_{n,\mathbf{s},j}^{(0)} : C_0(\mathbb{R}^n, K) \rightarrow W_{n,\mathbf{s},j}^{(0)}$ by

$$Q_{n,\mathbf{s},j}^{(0)} f = \sum_{\mathbf{k} \in \mathbb{Z}^n} \langle \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]}, f \rangle \psi_{\mathbf{s},j,\mathbf{k}}^{[n]}$$

for all $f \in C_0(\mathbb{R}^n, K)$ and $\mathbf{s} \in \{0, 1\}^n$.

We now have

$$V_{n,j}^{(0)} =_{\text{n.s.}} \bigotimes_{\varepsilon, k=1}^n V_{1,j}^{(0)},$$

$$W_{n,\mathbf{s},j}^{(0)} =_{\text{n.s.}} \bigotimes_{\varepsilon, k=1}^n W_{1,\mathbf{s}[k],j}^{(0)},$$

$$P_{n,j}^{(0)} = \bigotimes_{\varepsilon, k=1}^n P_{1,j}^{(0)}, \quad \text{and}$$

$$Q_{n,\mathbf{s},j}^{(0)} = \bigotimes_{\varepsilon, k=1}^n Q_{1,\mathbf{s}[k],j}^{(0)}.$$

We give now a general result on the tensor products of the function space $C_0(\mathbb{R}, K)$ with itself.

Theorem 5.1. *We have*

$$\bigotimes_{\varepsilon, j=1}^n C_0(\mathbb{R}, K) =_{\text{n.s.}} C_0(\mathbb{R}^n, K).$$

Proof. Define

$$F^{[n]} :=_{\text{n.s.}} \bigotimes_{\varepsilon}^n C_0(\mathbb{R}, K).$$

We have $F^{[n]} \subset C_0(\mathbb{R}^n, K)$. Let φ be the Deslauriers-Dubuc fundamental function of some degree m and $\varphi^{[n]}$ the n -dimensional tensor product mother scaling function generated by φ . Let spaces $V_{1,j}^{(0)}$, $j \in \mathbb{Z}$, belong to the interpolating multiresolution analysis generated by φ and spaces $V_{n,j}^{(0)}$, $j \in \mathbb{Z}$, be the corresponding tensor product spaces. Let $f \in C_0(\mathbb{R}^n, K)$. It follows from Theorem 4.3 that $P_{n,j}^{(0)}f \rightarrow f$ as $j \rightarrow \infty$. Now $P_{n,j}^{(0)}f \in V_{n,j}^{(0)} \subset F^{[n]}$ for all $j \in \mathbb{Z}$ from which it follows that $f \in \overline{F^{[n]}} = F^{[n]}$. Consequently $F^{[n]} =_{\text{n.s.}} C_0(\mathbb{R}^n, K)$. \square

For example, $C_0(\mathbb{R}) \hat{\otimes}_{\varepsilon} C_0(\mathbb{R}) =_{\text{n.s.}} C_0(\mathbb{R}^2)$. As in the case of $C_u(\mathbb{R}^n, K)$ we get

$$V_{n,j_0}^{(0)} \dot{+} \sum_{j=j_0}^{\infty} W_{n,j}^{(0)} \neq C_0(\mathbb{R}^n, K)$$

when the mother scaling function is Lipschitz continuous. We also have

$$C_0(\mathbb{R}^n, K) =_{\text{n.s.}} \text{clos} \left(\bigcup_{l=j_0}^{\infty} \left(V_{n,j_0}^{(0)} \dot{+} \sum_{j=j_0}^l W_{n,j}^{(0)} \right) \right)$$

for all $j_0 \in \mathbb{Z}$.

6. Interpolating Dual MRA

We shall have $K = \mathbb{R}$ or $K = \mathbb{C}$ and $n \in \mathbb{Z}_+$ throughout this section. We shall also have $E = C_u(\mathbb{R}^n, K)$ or $E = C_0(\mathbb{R}^n, K)$.

6.1. General

Definition 6.1. When $j \in \mathbb{Z}$ define

$$\begin{aligned} \tilde{V}_{n,j} &:=_{\text{n.s.}} \left\{ \sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{d}[\mathbf{k}] \tilde{\varphi}_{j,\mathbf{k}}^{[n]} : \mathbf{d} \in l^1(\mathbb{Z}^n, K) \right\} \\ \|\tilde{f}|_{\tilde{V}_{n,j}}\| &:= \|\tilde{f}|_{C_u(\mathbb{R}^n, K)^*}\| = \|\tilde{f}|_{C_0(\mathbb{R}^n, K)^*}\|, \quad \tilde{f} \in \tilde{V}_{n,j}. \end{aligned}$$

We may identify Banach space $\tilde{V}_{n,j}$ for both $E = C_u(\mathbb{R}^n, K)$ and $E = C_0(\mathbb{R}^n, K)$ for each $j \in \mathbb{Z}$. It follows also that function $\tilde{t}_{n,j} : l^1(\mathbb{Z}^n, K) \rightarrow \tilde{V}_{n,j}$ defined by

$$\tilde{t}_{n,j}(\mathbf{d}) := \sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{d}[\mathbf{k}] \tilde{\varphi}_{j,\mathbf{k}}^{[n]}$$

for all $\mathbf{d} \in l^1(\mathbb{Z}^n, K)$ is an isometric isomorphism from $l^1(\mathbb{Z}^n, K)$ onto $\tilde{V}_{n,j}$. We have

$$\tilde{f} \in \tilde{V}_{n,j} \iff \tilde{f}(2 \cdot) \in \tilde{V}_{n,j+1}$$

and

$$\tilde{f} \in \tilde{V}_{n,j} \iff \tilde{f}(\cdot - 2^{-j}\mathbf{k}) \in \tilde{V}_{n,j}.$$

for all $\tilde{f} \in C_u(\mathbb{R}^n, K)^*$, $j \in \mathbb{Z}$, and $\mathbf{k} \in \mathbb{Z}^n$.

Definition 6.2. When $j \in \mathbb{Z}$ and $\mathbf{s} \in \{0, 1\}^n$ define

$$\begin{aligned} \tilde{W}_{n,\mathbf{s},j} &:=_{\text{n.s.}} \left\{ \sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{d}[\mathbf{k}] \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]} : \mathbf{d} \in l^1(\mathbb{Z}^n, K) \right\} \\ \|\tilde{f}|_{\tilde{W}_{n,\mathbf{s},j}}\| &:= \|\tilde{f}|_{C_u(\mathbb{R}^n, K)^*}\| = \|\tilde{f}|_{C_0(\mathbb{R}^n, K)^*}\|, \quad \tilde{f} \in \tilde{W}_{n,\mathbf{s},j}. \end{aligned}$$

Spaces $\tilde{W}_{n,\mathbf{s},j}$, $\mathbf{s} \in \{0, 1\}^n$, $j \in \mathbb{Z}$, are topologically isomorphic to $l^1(\mathbb{Z}^n, K)$. Assume that $j \in \mathbb{Z}$, $\mathbf{d} \in l^1(J_+(n) \times \mathbb{Z}^n, K)$, and

$$\tilde{f} := \sum_{\mathbf{s} \in J_+(n)} \sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{d}[\mathbf{s}, \mathbf{k}] \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]}.$$

Now $\|\tilde{f}|_{C_u(\mathbb{R}^n, K)^*}\| = \|\tilde{f}|_{C_0(\mathbb{R}^n, K)^*}\|$. Consequently we may identify the Banach space $\tilde{W}_{n,\mathbf{s},j}$ for both $E = C_u(\mathbb{R}^n, K)$ and $E = C_0(\mathbb{R}^n, K)$ for each $\mathbf{s} \in \{0, 1\}^n$ and $j \in \mathbb{Z}$.

Definition 6.3. When $n \in \mathbb{Z}_+$ and $j \in \mathbb{Z}$ define

$$\tilde{P}_{n,j}\tilde{f} := \sum_{\mathbf{k} \in \mathbb{Z}^n} \langle \tilde{f}, \varphi_{j,\mathbf{k}}^{[n]} \rangle \tilde{\varphi}_{j,\mathbf{k}}^{[n]}$$

for all $\tilde{f} \in E^*$.

Definition 6.4. When $n \in \mathbb{Z}_+$, $\mathbf{s} \in \{0, 1\}^n$, and $j \in \mathbb{Z}$ define

$$\tilde{Q}_{n,\mathbf{s},j}\tilde{f} := \sum_{\mathbf{k} \in \mathbb{Z}^n} \langle \tilde{f}, \psi_{\mathbf{s},j,\mathbf{k}}^{[n]} \rangle \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]}$$

for all $\tilde{f} \in E^*$.

We have $\tilde{Q}_{n,\mathbf{0}_n,j} = \tilde{P}_{n,j}$ for all $j \in \mathbb{Z}$.

Lemma 6.1.

- (i) $\forall j \in \mathbb{Z}, \mathbf{s} \in \{0, 1\}^n, \mathbf{t} \in \{0, 1\}^n : \mathbf{s} \neq \mathbf{t} \implies \forall \tilde{f} \in \tilde{W}_{n,\mathbf{s},j} : \tilde{Q}_{n,\mathbf{t},j}\tilde{f} = 0$.
- (ii) Operator $\tilde{Q}_{n,\mathbf{s},j}$ is a continuous projection of E^* onto $\tilde{W}_{n,\mathbf{s},j}$ for all $j \in \mathbb{Z}$ and $\mathbf{s} \in \{0, 1\}^n$.
- (iii) $\tilde{Q}_{n,\mathbf{s},j} = \tilde{Q}_{n,\mathbf{s},j} \circ \tilde{P}_{n,j'}$ for all $j, j' \in \mathbb{Z}$, $j < j'$, and $\mathbf{s} \in \{0, 1\}^n$.

For example, when $n = 2$ and $\mathbf{t} = (0, 1)$ we have $\tilde{Q}_{n,\mathbf{t},j}\delta = 0$ for all $j \in \mathbb{Z}$. When $j \in \mathbb{Z}$, $\mathbf{s}, \mathbf{t} \in \{0, 1\}^n$, and $\mathbf{s} \neq \mathbf{t}$ we have $\tilde{W}_{n,\mathbf{s},j} \cap \tilde{W}_{n,\mathbf{t},j} = \{0\}$.

Definition 6.5. When $j \in \mathbb{Z}$ define

$$\tilde{W}_{n,j} :=_{\text{n.s.}} \sum_{\mathbf{s} \in J_+(n)} \tilde{W}_{n,\mathbf{s},j}.$$

Definition 6.6. When $j \in \mathbb{Z}$ define

$$\tilde{Q}_{n,j} := \sum_{\mathbf{s} \in J_+(n)} \tilde{Q}_{n,\mathbf{s},j}.$$

Assume that $j \in \mathbb{Z}$. Now operator $\tilde{Q}_{n,j}$ is a continuous projection of E^* onto $\tilde{W}_{n,j}$ and $\tilde{Q}_{n,j}\tilde{f} = 0$ for all $\tilde{f} \in \tilde{V}_{n,j}$. We also have $\tilde{Q}_{n,j} = \tilde{P}_{n,j+1} - \tilde{P}_{n,j}$ and $\tilde{V}_{n,j+1} =_{\text{n.s.}} \tilde{V}_{n,j} \dot{+} \tilde{W}_{n,j}$.

Theorem 6.1.

- (i) $\forall j \in \mathbb{Z}, \mathbf{s} \in \{0, 1\}^n, \mathbf{t} \in \{0, 1\}^n : \mathbf{s} \neq \mathbf{t} \iff \tilde{W}_{n,\mathbf{s},j} \perp W_{n,\mathbf{t},j}^{(u)}$
- (ii) $\forall j_1, j_2 \in \mathbb{Z} : j_1 \leq j_2 \iff \tilde{V}_{n,j_1} \perp W_{n,j_2}^{(u)}$
- (iii) $\forall j_1, j_2 \in \mathbb{Z} : j_1 \geq j_2 \iff \tilde{W}_{n,j_1} \perp V_{n,j_2}^{(u)}$
- (iv) $\forall j_1, j_2 \in \mathbb{Z} : j_1 \neq j_2 \iff \tilde{W}_{n,j_1} \perp W_{n,j_2}^{(u)}$

Proof. Let $j \in \mathbb{Z}, \mathbf{s}, \mathbf{t} \in \{0, 1\}^n$, and $\mathbf{k}, \ell \in \mathbb{Z}^n$. Proposition (i) follows from $\langle \tilde{\psi}_{\mathbf{s},j,\ell}^{[n]}, \psi_{\mathbf{t},j,\mathbf{k}}^{[n]} \rangle = \delta_{\mathbf{s},\mathbf{t}} \delta_{\ell,\mathbf{k}}$. Suppose that $j_1, j_2 \in \mathbb{Z}$ and $j_1 \neq j_2$. If $j_1 > j_2$ then $W_{n,j_2}^{(u)} \subset V_{n,j_1}^{(u)}$ and $\tilde{W}_{n,j_1} \perp V_{n,j_1}^{(u)}$ and hence $\tilde{W}_{n,j_1} \perp W_{n,j_2}^{(u)}$. If $j_1 < j_2$ then $\tilde{W}_{n,j_1} \subset \tilde{V}_{n,j_2}$ and $\tilde{V}_{n,j_2} \perp W_{n,j_2}^{(u)}$ and hence $\tilde{W}_{n,j_1} \perp W_{n,j_2}^{(u)}$. So proposition (iv) is true. \square

Theorem 6.2. Let $\tilde{f} \in C_0(\mathbb{R}^n, K)^*$. Then $\tilde{P}_{n,j}\tilde{f} \rightarrow \tilde{f}$ in the weak-* topology of $C_0(\mathbb{R}^n, K)^*$ as $j \rightarrow \infty$.

Proof. Let $\tilde{f} \in C_0(\mathbb{R}^n, K)^*$. Suppose that $f \in C_0(\mathbb{R}^n, K)$. Then

$$\begin{aligned} \langle \tilde{P}_{n,j}\tilde{f}, P_{n,j}^{(0)}f \rangle &= \sum_{\mathbf{k} \in \mathbb{Z}^n} \langle \tilde{f}, \varphi_{j,\mathbf{k}}^{[n]} \rangle \langle \tilde{\varphi}_{j,\mathbf{k}}^{[n]}, P_{n,j}^{(0)}f \rangle = \left\langle \tilde{f}, \sum_{\mathbf{k} \in \mathbb{Z}^n} \langle \tilde{\varphi}_{j,\mathbf{k}}^{[n]}, f \rangle \varphi_{j,\mathbf{k}}^{[n]} \right\rangle \\ &= \langle \tilde{f}, P_{n,j}^{(0)}f \rangle. \end{aligned}$$

There exists $c \in \mathbb{R}_+$ so that $\|\tilde{P}_{n,j}\| \leq c$ for all $j \in \mathbb{Z}$. Consequently

$$\begin{aligned} \left| \langle \tilde{f}, f \rangle - \langle \tilde{P}_{n,j}\tilde{f}, f \rangle \right| &\leq \left| \langle \tilde{f}, f \rangle - \langle \tilde{f}, P_{n,j}^{(0)}f \rangle \right| \\ &\quad + \left| \langle \tilde{P}_{n,j}\tilde{f}, P_{n,j}^{(0)}f \rangle - \langle \tilde{P}_{n,j}\tilde{f}, f \rangle \right| \\ &= \left| \langle \tilde{f}, f - P_{n,j}^{(0)}f \rangle \right| + \left| \langle \tilde{P}_{n,j}\tilde{f}, P_{n,j}^{(0)}f - f \rangle \right| \\ &\leq \|\tilde{f}\| \|f - P_{n,j}^{(0)}f\| + c \|\tilde{f}\| \|f - P_{n,j}^{(0)}f\| \rightarrow 0, \end{aligned}$$

as $j \rightarrow \infty$. \square

For example, let

$$\tilde{f}(x) := \begin{cases} x + 1; & x \in [-1, 0[\\ -x + 1; & x \in [0, 1[\\ 0; & \text{otherwise.} \end{cases}$$

and

$$f(x) := \begin{cases} -2x - 2; & x \in [-1, 0[\\ 2x - 2; & x \in [0, 1[\\ 0; & \text{otherwise.} \end{cases}$$

for all $x \in \mathbb{R}$. Now

$$\langle \tilde{P}_{1,j} \tilde{f}, f \rangle \rightarrow \langle \tilde{f}, f \rangle = \int_{\mathbb{R}} \tilde{f}(x) f(x) dx = -\frac{4}{3}$$

as $j \rightarrow \infty$.

6.2. Tensor Product Representation of the Dual MRA

It follows from Equation (2.1) that $C_0(\mathbb{R}^n, K)^* \hat{\otimes}_{\varepsilon^s} C_0(\mathbb{R}, K)^*$ is a closed subspace of $C_0(\mathbb{R}^{n+1}, K)^*$.

Definition 6.7. Let $j \in \mathbb{Z}$. Define

$$\tilde{M}_{n,j} :=_{\text{n.s.}} \bigotimes_{k=1}^n \hat{\otimes}_{(C_0(\mathbb{R}^k, K)^*)} \tilde{V}_{1,j}.$$

Banach space $\tilde{M}_{n,j}$ is a closed subspace of $C_0(\mathbb{R}^n, K)^*$.

Definition 6.8. Define function

$$\xi^{[n]} : \bigotimes_{k=1}^n \pi l^1(\mathbb{Z}, K) \rightarrow l^1(\mathbb{Z}^n, K)$$

by

$$\xi^{[n]}(\check{\mathbf{e}}_{\mathbf{k}}^{\otimes}) := \check{\mathbf{e}}_{\mathbf{k}}, \quad \mathbf{k} \in \mathbb{Z}^n,$$

and extending by linearity and continuity onto whole $\bigotimes_{k=1}^n \pi l^1(\mathbb{Z}, K)$.

Function $\xi^{[n]}$ is an isometric isomorphism from Banach space $\bigotimes_{k=1}^n \pi l^1(\mathbb{Z}, K)$ onto Banach space $l^1(\mathbb{Z}^n, K)$.

Lemma 6.2. When $j \in \mathbb{Z}$

$$\bigotimes_{k=1}^n \pi \tilde{V}_{1,j} =_{\text{n.s.}} \tilde{V}_{n,j}.$$

Proof. Let

$$E :=_{\text{n.s.}} \bigotimes_{k=1}^n \pi \tilde{V}_{1,j}$$

22

and

$$F :=_{\text{n.s.}} \bigotimes_{k=1}^n \pi l^1(\mathbb{Z}, K).$$

Let

$$\alpha := \bigotimes_{k=1}^n \tilde{l}_{1,j}$$

Function α is an isometric isomorphism from F onto E . Let $\beta := \alpha \circ (\xi^{[n]})^{-1}$. Now β is an isometric isomorphism from $l^1(\mathbb{Z}^n, K)$ onto E and $\beta(\tilde{\mathbf{e}}_{\mathbf{k}}) = \tilde{\varphi}_{j,\mathbf{k}}^{[n]} = \tilde{l}_{n,j}(\tilde{\mathbf{e}}_{\mathbf{k}})$ for all $\mathbf{k} \in \mathbb{Z}^n$. When $\mathbf{d} \in l^1(\mathbb{Z}^n, K)$ we have $\beta(\mathbf{d}) = \tilde{l}_{n,j}(\mathbf{d})$ and $\|\beta(\mathbf{d})\|_E = \|\tilde{l}_{n,j}(\mathbf{d})\|_{C_0(\mathbb{R}^n, K)^*}$. \square

Lemma 6.3. *When $j \in \mathbb{Z}$*

$$\tilde{V}_{n,j} =_{\text{n.s.}} \tilde{M}_{n,j} =_{\text{n.s.}} \bigotimes_{k=1}^n \tilde{V}_{1,j} =_{\text{n.s.}} \bigotimes_{k=1}^n \tilde{V}_{1,j}.$$

Proof. Use induction by n , metric approximation property of l^1 , and [41, prop. 7.1]. \square

Definition 6.9. *Define*

$$\tilde{N}_{n,\mathbf{s},j} :=_{\text{n.s.}} \bigotimes_{k=1}^n \tilde{W}_{1,\mathbf{s}[k],j}$$

where $j \in \mathbb{Z}$ and $\mathbf{s} \in \{0, 1\}^n$.

Definition 6.10. *Let $j \in \mathbb{Z}$ and $\mathbf{s} \in \{0, 1\}^n$. Define operator $\tilde{R}_{n,\mathbf{s},j} : \tilde{V}_{n,j+1} \rightarrow \tilde{N}_{n,\mathbf{s},j}$ by*

$$\tilde{R}_{n,\mathbf{s},j} = \bigotimes_{k=1}^n \left(\tilde{Q}_{1,\mathbf{s}[k],j} | \tilde{V}_{1,j+1} \right)$$

Assume that $j \in \mathbb{Z}$, and $\mathbf{s} \in \{0, 1\}^n$. Now

$$\tilde{R}_{n,\mathbf{s},j} \tilde{f} = \sum_{\mathbf{k} \in \mathbb{Z}^n} \langle \tilde{f}, \psi_{\mathbf{s},j,\mathbf{k}}^{[n]} \rangle \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]} \quad (6.1)$$

for all $\tilde{f} \in \tilde{V}_{n,j+1}$. The series in Equation (6.1) converges absolutely for all $\tilde{f} \in \tilde{V}_{n,j+1}$. We also have $\tilde{W}_{n,\mathbf{s},j} =_{\text{n.s.}} \tilde{N}_{n,\mathbf{s},j}$ and $\tilde{Q}_{n,\mathbf{s},j} = \tilde{R}_{n,\mathbf{s},j} \circ \tilde{P}_{n,j+1}$.

7. Besov Space Norm Equivalence

7.1. Norm Equivalence for the Besov Spaces in the n -dimensional Case

This derivation is based on the corresponding one-dimensional derivation in [17]. The cases $p < 1$ or $q < 1$ yielding quasi-Banach spaces $B_{p,q}^\sigma$ are not discussed in this article. We assume that $n \in \mathbb{Z}_+$ throughout this section.

We give first some definitions similar to those in [38] related to orthonormal wavelets.

Definition 7.1. Let $\bar{\varphi} : \mathbb{R}^n \rightarrow K$ be a mother scaling function of an orthonormal wavelet family. When $j \in \mathbb{Z}$, $\mathbf{k} \in \mathbb{Z}^n$, and $f \in L^\infty(\mathbb{R}^n)$ define

$$\bar{\alpha}_{j,\mathbf{k}}(f) := 2^{nj} \int_{\mathbf{x} \in \mathbb{R}^n} \bar{\varphi}^*(2^j \mathbf{x} - \mathbf{k}) f(\mathbf{x}) d\tau.$$

The spaces $V_j(p)$ that are defined in [38] are denoted by $\bar{V}_j(p)$ in this document. Space $\bar{V}_j(p)$ is a closed subspace of $L^p(\mathbb{R}^n)$ for each $j \in \mathbb{Z}$ and $p \in [1, \infty]$ at least when the mother scaling function $\bar{\varphi}$ is continuous and compactly supported.

Definition 7.2. Let $\bar{\varphi} : \mathbb{R}^n \rightarrow \mathbb{C}$ be a scaling function of an orthonormal wavelet family. Let $p \in [1, \infty]$ and $j \in \mathbb{Z}$. Define operator $\bar{P}_j^{(p)} : L^p(\mathbb{R}^n) \rightarrow \bar{V}_j(p)$ by

$$(\bar{P}_j^{(p)} f)(\mathbf{x}) := \sum_{\mathbf{k} \in \mathbb{Z}^n} \bar{\alpha}_{j,\mathbf{k}}(f) \bar{\varphi}(2^j \mathbf{x} - \mathbf{k})$$

for all $f \in L^p(\mathbb{R}^n)$ and $\mathbf{x} \in \mathbb{R}^n$. Define also $\bar{Q}_j^{(p)} := \bar{P}_{j+1}^{(p)} - \bar{P}_j^{(p)}$.

When $\bar{\varphi}$ is a compactly supported and continuous function operator $\bar{P}_j^{(p)}$ is a continuous linear projection of $L^p(\mathbb{R}^n)$ onto $\bar{V}_j(p)$ for each $j \in \mathbb{Z}$ and $p \in [1, \infty]$.

Definition 7.3. Let $p \in [1, \infty]$, $q \in [1, \infty]$, and $\sigma \in \mathbb{R}_+$. Let $j_0 \in \mathbb{Z}$ and $\bar{r} \in \mathbb{Z}_+$, $\bar{r} > \sigma$. Let $\bar{\varphi} : \mathbb{R}^n \rightarrow \mathbb{C}$ be a mother scaling function of an \bar{r} -regular orthonormal MRA of $L^2(\mathbb{R}^n)$ and $\bar{P}_j^{(p)}$ and $\bar{Q}_j^{(p)}$, $j \in \mathbb{Z}$, be the corresponding projection operators. When $f \in L^p(\mathbb{R}^n)$ define

$$\|f\|_{B_{p,q}^{\sigma}(\mathbb{R}^n);j_0}^{(o)} := \left\| \bar{P}_{j_0}^{(p)} f \right\|_{L^p(\mathbb{R}^n)} + \|\bar{\mathbf{h}}\|_{l^q(\mathbb{N} + j_0)}$$

where

$$\begin{aligned} \bar{h}_j &:= 2^{j\sigma} \left\| \bar{Q}_j^{(p)} f \right\|_{L^p(\mathbb{R}^n)}, \quad j \in \mathbb{N} + j_0 \\ \bar{\mathbf{h}} &:= (\bar{h}_j)_{j=j_0}^\infty. \end{aligned}$$

Norm $\|\cdot\|_{B_{p,q}^{\sigma}(\mathbb{R}^n);j_0}^{(o)}$ is an equivalent norm for the Besov space $B_{p,q}^\sigma(\mathbb{R}^n)$ and characterizes $B_{p,q}^\sigma(\mathbb{R}^n)$ on $\text{bor}(\mathbb{R}^n, \mathbb{C})$. This has been proved in [38, chapter 2.9 proposition 4].

Definition 7.4. Let $p \in [1, \infty]$, $q \in [1, \infty]$, and $\sigma \in \mathbb{R}_+$. Let $j_0 \in \mathbb{Z}$. Let $P_{n,j}^{(u)}$ and $Q_{n,j}^{(u)}$, $j \in \mathbb{Z}$, be the projection operators belonging to a compactly supported interpolating tensor product MRA of $C_u(\mathbb{R}^n)$. When $f \in C_u(\mathbb{R}^n)$ define

$$\|f\|_{B_{p,q}^{(\sigma)}(\mathbb{R}^n);j_0}^{(i)} := \left\| P_{n,j_0}^{(u)} f |L^p(\mathbb{R}^n)\right\| + \|\mathbf{h}\|_{l^q(\mathbb{N} + j_0)}$$

where

$$h_j := 2^{j\sigma} \left\| Q_{n,j}^{(u)} f |L^p(\mathbb{R}^n)\right\|, \quad j \in \mathbb{N} + j_0$$

$$\mathbf{h} := (h_j)_{j=j_0}^\infty.$$

Definition 7.5. Let $p \in [1, \infty]$, $q \in [1, \infty]$, and $\sigma \in \mathbb{R}_+$. Let $j_0 \in \mathbb{Z}$. Let $\tilde{\varphi}_{j,\mathbf{k}}^{[n]}$ and $\tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]}$, $j \in \mathbb{Z}$, $\mathbf{s} \in J_+(n)$, $\mathbf{k} \in \mathbb{Z}^n$, be the dual scaling functions and dual wavelets belonging to a compactly supported interpolating tensor product MRA of $C_u(\mathbb{R}^n)$. When $f \in C_u(\mathbb{R}^n)$ define

$$\|f\|_{B_{p,q}^{(w)}(\mathbb{R}^n);j_0}^{(w)} := \left\| \left(\left\langle \tilde{\varphi}_{j_0,\mathbf{k}}^{[n]}, f \right\rangle \right)_{\mathbf{k} \in \mathbb{Z}^n} \right\|_p$$

$$+ \left\| \left(2^{(\sigma - \frac{n}{p})j} \left\| \left(\left\langle \tilde{\psi}_{\mathbf{s},j,\ell}^{[n]}, f \right\rangle \right)_{\mathbf{s} \in J_+(n), \ell \in \mathbb{Z}^n} \right\|_p \right)_{j=j_0}^\infty \right\|_q.$$

Definition 7.6. When $j \in \mathbb{Z}$ and $p \in [1, \infty]$ define

$$V_{n,j}^{(u)}(p) := \left\{ \mathbf{x} \in \mathbb{R}^n \mapsto \sum_{\mathbf{k} \in \mathbb{Z}^n} \mathbf{a}[\mathbf{k}] \varphi_{j,\mathbf{k}}^{[n]}(\mathbf{x}) : \mathbf{a} \in l^p(\mathbb{Z}^n) \right\}$$

and $\|f|V_{n,j}^{(u)}(p)\| := \|f|L^p(\mathbb{R}^n)\|$ for all $f \in V_{n,j}^{(u)}(p)$.

Definition 7.7. Let $j \in \mathbb{Z}$ and $p \in [1, \infty]$. When $T \in \mathcal{L}(L^p(\mathbb{R}^n), L^p(\mathbb{R}^n))$ define

$$\|T\|_{V_{n,j}^{(u)}(p)} := \left\| (T|_{V_{n,j}^{(u)}(p)}) | \mathcal{L}(V_{n,j}^{(u)}(p), L^p(\mathbb{R}^n)) \right\|.$$

Definition 7.8. Let $j \in \mathbb{Z}$ and $p \in [1, \infty]$. When $T \in \mathcal{L}(C_u(\mathbb{R}^n), C_u(\mathbb{R}^n))$ and $T[\bar{V}_j(p)] \subset L^p(\mathbb{R}^n)$ define

$$\|T\|_{\bar{V}_j(p)} := \left\| (T|_{\bar{V}_j(p)}) | \mathcal{L}(\bar{V}_j(p), L^p(\mathbb{R}^n)) \right\|.$$

Definition 7.9. When $p \in [1, \infty]$ define

$$c_{\text{interp}}(p) := \begin{cases} \frac{p}{p-1}; & p \in]1, \infty[\\ 1; & p = 1 \vee p = \infty. \end{cases}$$

Lemma 7.1. Suppose that $q \in [1, \infty]$ and $t \in \mathbb{R}_+$. Let $a(j, j') := 2^{-|j'-j|t}$ for all $j, j' \in \mathbb{N}$. Define

$$Ab := \left(\sum_{j'=0}^\infty a(j, j') b_{j'} \right)_{j \in \mathbb{N}}$$

for all $\mathbf{b} \in l^q(\mathbb{N})$. Then $A \in \mathcal{L}(l^q(\mathbb{N}), l^q(\mathbb{N}))$.

Proof. Cases $q = 1$ and $q = \infty$ can be proved by starting from the expression of $\mathbf{A}\mathbf{b}$ where $\mathbf{b} \in l^q(\mathbb{N})$. When $q \in]1, \infty[$ the proof is based on interpolation of Banach spaces, see [2]. \square

Theorem 7.1. *Let $p \in [1, \infty]$, $q \in [1, \infty]$, $j_0 \in \mathbb{Z}$, $\sigma \in \mathbb{R}_+$, and $n/p < \sigma < r_0$. Let $\varphi^{[n]}$ be a scaling function of a compactly supported interpolating tensor product MRA of $C_u(\mathbb{R}^n)$ and $\bar{\varphi} : \mathbb{R}^n \rightarrow \mathbb{C}$ a compactly supported and continuous mother scaling function of a $([\sigma] + 1)$ -regular orthonormal wavelet family. Suppose that $\varphi^{[n]} \in C^{r_0}(\mathbb{R}^n)$. Suppose also that $(\tilde{\varphi}^{[n]}, \varphi^{[n]})$ and $(2^{nj}\bar{\varphi}^*, \bar{\varphi})$ span all polynomials of degree at most $[\sigma] - 1$. Let B be the normed space $B_{p,q}^\sigma(\mathbb{R}^n) \cap C_u(\mathbb{R}^n)$ equipped with some Besov space norm. Then $\|\cdot\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(i)}$ and $\|\cdot\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(o)}$ are equivalent norms on the vector space B and norm $\|\cdot\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(i)}$ characterizes B on $C_u(\mathbb{R}^n)$.*

Proof. Let $\bar{r}_0 := [\sigma] + 1$. We prove first that $\|f\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(i)} \leq c\|f\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(o)}$ for all $f \in B$ and for some $c \in \mathbb{R}_+$. Define q' by

$$\frac{1}{q'} + \frac{1}{q} = 1.$$

There exists $c_1 \in \mathbb{R}_+$ so that

$$\left\| P_{n,j}^{(u)} \right\|_{\bar{V}_{j'}(p)} \leq c_1 \cdot 2^{n(j'-j)/p} \quad (7.1)$$

for all $j, j' \in \mathbb{Z}$, $j' \geq j$. There exists $c_2 \in \mathbb{R}_+$ so that

$$\left\| I - P_{n,j}^{(u)} \right\|_{\bar{V}_{j'}(p)} \leq c_2 \cdot c_{\text{interp}}(p) \cdot 2^{(j'-j)\bar{r}_0} \quad (7.2)$$

for all $j, j' \in \mathbb{Z}$, $j' \leq j$. These two results can be proved by Banach space interpolation between cases $p = 1$ and $p = \infty$. Let $f \in B$. We have

$$\begin{aligned} \left\| P_{n,j_0}^{(u)} f \right\|_p &= \left\| P_{n,j_0}^{(u)} \left(\bar{P}_{j_0}^{(p)} f + \sum_{j=j_0}^{\infty} \bar{Q}_j^{(p)} f \right) \right\|_p \\ &\leq \left\| P_{n,j_0}^{(u)} \right\|_{\bar{V}_{j_0}(p)} \left\| \bar{P}_{j_0}^{(p)} f \right\|_p + \sum_{j=j_0}^{\infty} \left\| P_{n,j_0}^{(u)} \right\|_{\bar{V}_{j+1}(p)} \left\| \bar{Q}_j^{(p)} f \right\|_p \\ &\leq c_3 \left\| \bar{P}_{j_0}^{(p)} f \right\|_p + c_3 \cdot 2^{(n-nj_0)/p} \sum_{j=j_0}^{\infty} 2^{-j(\sigma - \frac{n}{p})} \bar{\mathbf{h}}[j] \end{aligned}$$

where $\bar{\mathbf{h}} := (2^{j\sigma} \|\bar{Q}_j^{(p)} f\|_p)_{j=j_0}^{\infty}$. Let

$$c_4 := c_3 \cdot 2^{(n-nj_0)/p} \cdot \left\| \left(2^{-j'(\sigma - \frac{n}{p})} \right)_{j'=j_0}^{\infty} \right\|_{q'}.$$

26

Now

$$\left\| P_{n,j_0}^{(u)} f \right\|_p \leq c_3 \left\| \bar{P}_{j_0}^{(p)} f \right\|_p + c_4 \|\bar{\mathbf{h}}\|_q \leq \max\{c_3, c_4\} \cdot \|f\|_{B_{p,q}^{\sigma}(\mathbb{R}^n); j_0}^{(o)}$$

where c_3 and c_4 do not depend on f . We have

$$\begin{aligned} \left\| Q_{n,j}^{(u)} f \right\|_p &= \left\| Q_{n,j}^{(u)} \left(\bar{P}_{j_0}^{(p)} f + \sum_{j'=j_0}^{\infty} \bar{Q}_{j'}^{(p)} f \right) \right\|_p \\ &\leq \left\| Q_{n,j}^{(u)} \right\|_{\bar{V}_{j_0}(p)} \left\| \bar{P}_{j_0}^{(p)} f \right\|_p + \sum_{j'=j_0}^{\infty} \left\| Q_{n,j}^{(u)} \right\|_{\bar{V}_{j'+1}(p)} \left\| \bar{Q}_{j'}^{(p)} f \right\|_p \end{aligned}$$

for all $j \in \mathbb{Z}$, $j \geq j_0$. It follows from Equation (7.2) that

$$2^{j\sigma} \left\| Q_{n,j}^{(u)} \right\|_{\bar{V}_{j_0}(p)} \left\| \bar{P}_{j_0}^{(p)} f \right\|_p \leq c_5 \cdot 2^{-j(\bar{r}_0 - \sigma)} \left\| \bar{P}_{j_0}^{(p)} f \right\|_p$$

for all $j \in \mathbb{Z}$, $j \geq j_0$, where c_5 is independent of f . Consequently

$$\begin{aligned} \left\| \left(2^{j\sigma} \left\| Q_{n,j}^{(u)} \right\|_{\bar{V}_{j_0}(p)} \left\| \bar{P}_{j_0}^{(p)} f \right\|_p \right)_{j=j_0}^{\infty} \right\|_q &\leq c_5 \left\| \left(2^{-j(\bar{r}_0 - \sigma)} \right)_{j=j_0}^{\infty} \right\|_q \left\| \bar{P}_{j_0}^{(p)} f \right\|_p \\ &= c_6 \left\| \bar{P}_{j_0}^{(p)} f \right\|_p \leq c_6 \|f\|_{B_{p,q}^{\sigma}(\mathbb{R}^n); j_0}^{(o)}. \end{aligned} \quad (7.3)$$

When $j' < j$ it follows from Equation (7.2) that

$$2^{j\sigma} \left\| Q_{n,j}^{(u)} \right\|_{\bar{V}_{j'+1}(p)} \left\| \bar{Q}_{j'}^{(p)} f \right\|_p \leq c_7 \cdot 2^{-|j'-j|(\bar{r}_0 - \sigma)} \bar{\mathbf{h}}[j'] \quad (7.4)$$

When $j' \geq j$ it follows from Equation (7.1) that

$$2^{j\sigma} \left\| Q_{n,j}^{(u)} \right\|_{\bar{V}_{j'+1}(p)} \left\| \bar{Q}_{j'}^{(p)} f \right\|_p \leq c_8 \cdot 2^{-|j'-j|(\sigma - \frac{n}{p})} \bar{\mathbf{h}}[j']. \quad (7.5)$$

Let

$$w := \min \left\{ \sigma - \frac{n}{p}, \bar{r}_0 - \sigma \right\}$$

and $c_9 := \max\{c_7, c_8\}$. Now by Equations (7.4) and (7.5)

$$\begin{aligned} a &:= \left\| \left(2^{j\sigma} \sum_{j'=j_0}^{\infty} \left\| Q_{n,j}^{(u)} \right\|_{\bar{V}_{j'+1}(p)} \left\| \bar{Q}_{j'}^{(p)} f \right\|_p \right)_{j=j_0}^{\infty} \right\|_q \\ &\leq c_9 \left\| \left(\sum_{j'=j_0}^{\infty} 2^{-|j'-j|w} \bar{\mathbf{h}}[j'] \right)_{j=j_0}^{\infty} \right\|_q. \end{aligned}$$

It follows from Lemma 7.1 that

$$a \leq c_{10} \|\bar{\mathbf{h}}\|_q \leq c_{10} \|f\|_{B_{p,q}^{\sigma}(\mathbb{R}^n); j_0}^{(o)}. \quad (7.6)$$

where c_{10} depends only on w and j_0 . By Equations (7.3) and (7.6) we have $\|f\|_{B_{p,q}^{\sigma,(\mathbb{R}^n)};j_0}^{(i)} \leq c_{10} \|f\|_{B_{p,q}^{\sigma,(\mathbb{R}^n)};j_0}^{(o)}$

We prove then that $\|g\|_{B_{p,q}^{\sigma,(\mathbb{R}^n)};j_0}^{(o)} \leq c \|g\|_{B_{p,q}^{\sigma,(\mathbb{R}^n)};j_0}^{(i)}$ for all $g \in B$ and for some $c \in \mathbb{R}_+$. Now

$$\begin{aligned} \left\| P_{n,j_0}^{(u)} g \right\|_p + \sum_{j=j_0}^{\infty} \left\| Q_{n,j}^{(u)} g \right\|_p &= \left\| P_{n,j_0}^{(u)} g \right\|_p + \sum_{j=j_0}^{\infty} \left\| 2^{-j\sigma} \cdot 2^{j\sigma} Q_{n,j}^{(u)} g \right\|_p \\ &\leq \left\| P_{n,j_0}^{(u)} g \right\|_p + \left\| \left(2^{j\sigma} \left\| Q_{n,j}^{(u)} g \right\|_p \right)_{j=j_0}^{\infty} \right\|_q \left\| \sum_{j=j_0}^{\infty} 2^{-j\sigma} \right\|_p \\ &\leq c_{11} \|g\|_{B_{p,q}^{\sigma,(\mathbb{R}^n)};j_0}^{(i)} \leq c_{12} \|g\|_{B_{p,q}^{\sigma,(\mathbb{R}^n)};j_0}^{(o)} \in \mathbb{R}_0 \end{aligned}$$

where c_{11} and c_{12} do not depend on g . Consequently

$$P_{n,j_0}^{(u)} g + \sum_{j=j_0}^m Q_{n,j}^{(u)} g \rightarrow g$$

in $L^p(\mathbb{R}^n)$ as $m \rightarrow \infty$. We have

$$\begin{aligned} \left\| \bar{P}_{j_0}^{(p)} g \right\|_p &= \left\| \bar{P}_{j_0}^{(p)} \left(P_{n,j_0}^{(u)} g + \sum_{j=j_0}^{\infty} Q_{n,j}^{(u)} g \right) \right\|_p \\ &\leq \left\| \bar{P}_{j_0}^{(p)} \right\| \left\| P_{n,j_0}^{(u)} g \right\|_p + \sum_{j'=j_0}^{\infty} \left\| \bar{P}_{j_0}^{(p)} \right\| \left\| Q_{n,j'}^{(u)} g \right\|_p \\ &\leq c_{13} \left\| P_{n,j_0}^{(u)} g \right\|_p + c_{13} \sum_{j=j_0}^{\infty} 2^{-j\sigma} \mathbf{h}[j] \\ &\leq c_{13} \left\| P_{n,j_0}^{(u)} g \right\|_p + c_{13} \left\| (2^{-j\sigma})_{j=j_0}^{\infty} \right\|_{q'} \|\mathbf{h}\|_q \\ &\leq c_{14} \|g\|_{B_{p,q}^{\sigma,(\mathbb{R}^n)};j_0}^{(i)} \end{aligned}$$

where $\mathbf{h} := (2^{j\sigma} \|Q_{n,j}^{(u)} g\|_p)_{j=j_0}^{\infty}$. There exists $c_{15} \in \mathbb{R}_+$ so that

$$\left\| I - \bar{P}_j^{(p)} \right\|_{V_{n,j'}^{(u)}(p)} \leq c_{15} \cdot c_{\text{interp}}(p) \cdot 2^{(j'-j)\sigma} \quad (7.7)$$

for all $j, j' \in \mathbb{Z}$, $j' \leq j$. This can be proved by Banach space interpolation between cases $p = 1$ and $p = \infty$. Furthermore,

$$\begin{aligned} \left\| \bar{Q}_j^{(p)} g \right\|_p &= \left\| \bar{Q}_j^{(p)} \left(P_{n,j_0}^{(u)} g + \sum_{j'=j_0}^{\infty} Q_{n,j'}^{(u)} g \right) \right\|_p \\ &\leq \left\| \bar{Q}_j^{(p)} \right\| \left\| P_{n,j_0}^{(u)} g \right\|_p + \sum_{j'=j_0}^{\infty} \left\| \bar{Q}_j^{(p)} \right\| \left\| Q_{n,j'}^{(u)} g \right\|_p. \end{aligned}$$

It follows from Equation (7.7) that

$$2^{j\sigma} \left\| \bar{Q}_j^{(p)} \right\| \left\| P_{n,j_0}^{(u)} g \right\|_p \leq c_{16} \cdot 2^{-j(r_0-\sigma)} \left\| P_{n,j_0}^{(u)} g \right\|_p$$

for all $j \geq j_0$ where c_{16} is independent of g . Consequently

$$\begin{aligned} \left\| \left(2^{j\sigma} \left\| \bar{Q}_j^{(p)} \right\| \left\| P_{n,j_0}^{(u)} g \right\|_p \right)_{j=j_0}^{\infty} \right\|_q &\leq c_{16} \left\| \left(2^{-j(r_0-\sigma)} \right)_{j=j_0}^{\infty} \right\|_q \left\| P_{n,j_0}^{(u)} g \right\|_p \\ &= c_{17} \left\| P_{n,j_0}^{(u)} g \right\|_p \leq c_{17} \|g\|_{B_{p,q}^{\sigma}(\mathbb{R}^n);j_0}. \end{aligned} \quad (7.8)$$

When $j' < j$ it follows from Equation (7.7) that

$$2^{j\sigma} \left\| \bar{Q}_j^{(p)} \right\| \left\| Q_{n,j'}^{(u)} g \right\|_p \leq c_{18} \cdot 2^{-|j'-j|(r_0-\sigma)} \mathbf{h}[j']. \quad (7.9)$$

When $j' \geq j$ it follows from uniform boundedness of the projection operators that

$$2^{j\sigma} \left\| \bar{Q}_j^{(p)} \right\| \left\| Q_{n,j'}^{(u)} g \right\|_p \leq c_{19} \cdot 2^{-|j'-j|\sigma} \mathbf{h}[j']. \quad (7.10)$$

Let

$$z := \min \{ \sigma, r_0 - \sigma \}.$$

Now by Equations (7.9) and (7.10) we have

$$\begin{aligned} b &:= \left\| \left(\sum_{j'=j_0}^{\infty} 2^{j\sigma} \left\| \bar{Q}_j^{(p)} \right\| \left\| Q_{n,j'}^{(u)} g \right\|_p \right)_{j=j_0}^{\infty} \right\|_q \\ &\leq c_{20} \left\| \left(\sum_{j'=j_0}^{\infty} 2^{-|j'-j|z} \mathbf{h}[j'] \right)_{j=j_0}^{\infty} \right\|_q. \end{aligned}$$

It follows from Lemma 7.1 that

$$b \leq c_{21} \|\mathbf{h}\|_q \leq c_{21} \|f\|_{B_{p,q}^{\sigma}(\mathbb{R}^n);j_0}. \quad (7.11)$$

where c_{21} depends only on z and j_0 . By Equations (7.8) and (7.11) we have $\|g\|_{B_{p,q}^{(o)}(\mathbb{R}^n);j_0} \leq c_{22} \|g\|_{B_{p,q}^{(i)}(\mathbb{R}^n);j_0}$.

Suppose then that $\|g\|_{B_{p,q}^{(o)}(\mathbb{R}^n);j_0} = \infty$. Let

$$\mathbf{A}b := \left(\sum_{j'=j_0}^{\infty} 2^{-|j'-j|z} \mathbf{b}[j'] \right)_{j=j_0}^{\infty}, \quad \mathbf{b} \in \mathbb{C}^{\mathbb{N}+j_0}.$$

Now

$$\infty = \left\| \left(2^{j\sigma} \left\| \bar{Q}_j^{(p)} g \right\|_p \right)_{j=j_0}^{\infty} \right\|_q \leq c_{17} \left\| P_{n,j_0}^{(u)} g \right\|_p + c_{20} \|\mathbf{A}b\|_q.$$

By the continuity of $A|_{l^q(\mathbb{N}+j_0)}$ we must have $\|\mathbf{h}\|_q = \infty$. Consequently $\|g\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(i)} = \infty$. \square

It is possible to construct an orthonormal compactly supported Daubechies wavelet family with arbitrary high regularity $\bar{r} \in \mathbb{R}_+$ and polynomial span (number of vanishing moments) $\bar{d} \in \mathbb{N}$ [9]. Consequently, when $j_0 \in \mathbb{Z}$, $\sigma \in \mathbb{R}_+$, $p \in [1, \infty]$, $q \in [1, \infty]$, $n/p < \sigma < r_0$, $\varphi^{[n]} \in C^{r_0}(\mathbb{R}^n)$, and $(\tilde{\varphi}^{[n]}, \varphi^{[n]})$ spans all polynomials of degree at most $\lceil \sigma \rceil - 1$ function $\varphi^{[n]}$ defines an equivalent norm $\|\cdot\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(i)}$ on Banach space B .

Theorem 7.2. *Let $p \in [1, \infty]$, $q \in [1, \infty]$, $\sigma \in \mathbb{R}_+$, $j_0 \in \mathbb{Z}$, $r_0 \in \mathbb{R}_+$, and $n/p < \sigma < r_0$. Let $\varphi^{[n]}$ be a scaling function of a compactly supported interpolating tensor product MRA of $C_u(\mathbb{R}^n)$ and suppose that $\varphi^{[n]} \in C^{r_0}(\mathbb{R}^n)$. Let B be the normed space $B_{p,q}^\sigma(\mathbb{R}^n) \cap C_u(\mathbb{R}^n)$ equipped with some Besov space norm. Then $\|\cdot\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(i)}$ and $\|\cdot\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(w)}$ are equivalent to the norm of B and they characterize B on $C_u(\mathbb{R}^n)$.*

Proof. Define

$$\mathbf{a}_j(f) := \left(\left\langle \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]}, f \right\rangle \right)_{\mathbf{s} \in J_+(n), \mathbf{k} \in \mathbb{Z}^n}$$

where $f \in B$ and $j \in \mathbb{Z}$. By applying Banach space interpolation to cases $p = 1$ and $p = \infty$ we get

$$\begin{aligned} c_1 \cdot 2^{-\frac{nj_0}{p}} \left\| \left(\left\langle \tilde{\varphi}_{j_0,\mathbf{k}}^{[n]}, f \right\rangle \right)_{\mathbf{k} \in \mathbb{Z}^n} \right\|_p &\leq \|P_{n,j_0}^{(u)} f\| \\ &\leq c_2 \cdot 2^{-\frac{nj_0}{p}} \left\| \left(\left\langle \tilde{\varphi}_{j_0,\mathbf{k}}^{[n]}, f \right\rangle \right)_{\mathbf{k} \in \mathbb{Z}^n} \right\|_p \end{aligned} \quad (7.12)$$

and

$$c_3 \cdot 2^{-\frac{nj}{p}} \|\mathbf{a}_j(f)\|_p \leq \|Q_{n,j}^{(u)} f\|_p \leq c_4 \cdot 2^{-\frac{nj}{p}} \|\mathbf{a}_j(f)\|_p$$

for all $j \in \mathbb{N} + j_0$ and $f \in B$. Consequently

$$\begin{aligned} c_3 \left\| \left(2^{j(\sigma - \frac{n}{p})} \|\mathbf{a}_j(f)\|_p \right)_{j=j_0}^\infty \right\|_q &\leq \left\| \left(2^{j\sigma} \|Q_{n,j}^{(u)} f\|_p \right)_{j=j_0}^\infty \right\|_q \\ &\leq c_4 \left\| \left(2^{j(\sigma - \frac{n}{p})} \|\mathbf{a}_j(f)\|_p \right)_{j=j_0}^\infty \right\|_q \end{aligned} \quad (7.13)$$

for all $f \in B$.

Let $c_5 := \min\{c_1 \cdot 2^{-nj_0/p}, c_3\}$ and $c_6 := \max\{c_2 \cdot 2^{-nj_0/p}, c_4\}$. It follows from Equations (7.12) and (7.13) that

$$c_5 \|f\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(w)} \leq \|f\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(i)} \leq c_6 \|f\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(w)}$$

for all $f \in B$. \square

It is possible to construct Deslauriers-Dubuc wavelet families with arbitrary high regularity $r \in \mathbb{R}_+$ (i.e. $\varphi^{[n]} \in C^r(\mathbb{R}^n)$) and polynomial span $d \in \mathbb{N}$ [17]. As a direct consequence of Theorems 7.1 and 7.2 norms $\|\cdot\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(w)}$ and $\|\cdot\|_{B_{p,q}^\sigma(\mathbb{R}^n);j_0}^{(i)}$ are equivalent to the restriction of some norm of Besov space $B_{p,q}^\sigma(\mathbb{R}^n)$ onto the vector space $B_{p,q}^\sigma(\mathbb{R}^n) \cap C_u(\mathbb{R}^n)$ and these norms characterize $B_{p,q}^\sigma(\mathbb{R}^n) \cap C_u(\mathbb{R}^n)$ on $C_u(\mathbb{R}^n)$.

7.2. Consequences of the Besov Space Norm Equivalence

In particular, the Besov space norm equivalence holds for Hölder spaces $C^\sigma(\mathbb{R}^n)$ for $\sigma \in \mathbb{R}_+ \setminus \mathbb{Z}$. When $n \in \mathbb{Z}_+$, $\sigma \in \mathbb{R}_+$, and $q \in [1, \infty]$ we have $B_{\infty,q}^\sigma(\mathbb{R}^n) \subset C_u(\mathbb{R}^n)$ [44, prop. 2.3.2/2(i) and eq. (2.3.5/1)].

Definition 7.10. When $n \in \mathbb{Z}_+$ and $j_0 \in \mathbb{Z}$ define

$$\Omega(n, j_0) := \{(\mathbf{0}_n, j_0, \mathbf{k}) : \mathbf{k} \in \mathbb{Z}^n\} \cup \bigcup_{j=j_0}^{\infty} \bigcup_{\mathbf{s} \in J_+(n)} \{(\mathbf{s}, j, \mathbf{k}) : \mathbf{k} \in \mathbb{Z}^n\}.$$

As the Banach space $B_{\infty,\infty}^\sigma(\mathbb{R}^n) =_{\text{tvs}} \mathcal{Z}^\sigma(\mathbb{R}^n)$ is isomorphic to l^∞ [44, section 2.5.5. page 87] the set $\{\psi_\alpha : \alpha \in \Omega(n, j_0)\}$ cannot be a Schauder basis of $B_{\infty,\infty}^\sigma(\mathbb{R}^n)$ with any summing order.

Remark 7.1. Suppose that $n \in \mathbb{Z}_+$, $j_0 \in \mathbb{Z}$, $\sigma \in \mathbb{R}_+$, and $\varphi^{[n]} \in C^r(\mathbb{R}^n)$ where $r \in \mathbb{R}_+$, $r > \sigma$. There exists $r_1 \in \mathbb{Z}_+$ so that $\text{supp } \varphi^{[n]} \in \overline{B}_{\mathbb{R}^n}(0; r_1)$. Consider

$$f := \sum_{k=0}^{\infty} 2^{-(j_0+k)\sigma} \psi_{\mathbf{e}_1^{[n]}, j_0+k, \eta(k)}^{[n]}$$

where

$$\begin{aligned} c &:= 2r_1 + \lceil 2^{j_0} \rceil (2\lceil \sigma \rceil + 1) \\ \eta(k) &:= c \cdot 2^k k \mathbf{e}_1^{[n]} \end{aligned}$$

for all $k \in \mathbb{Z}$, $k \geq j_0$, and the series converges in $C_0(\mathbb{R}^n)$. Let $m := \lceil r \rceil$ and

$$g_k := 2^{-(j_0+k)\sigma} \psi_{\mathbf{e}_1^{[n]}, j_0+k, \eta(k)}^{[n]}$$

for all $k \in \mathbb{N}$. As $\psi_{\mathbf{e}_1^{[n]}}^{[n]} \in B_{\infty,\infty}^\sigma(\mathbb{R}^n)$ it follows that there exists $c_1 \in \mathbb{R}_+$ so that

$$\omega_\infty^m(g_k; t) \leq c_1 t^\sigma \quad (7.14)$$

for all $k \in \mathbb{N}$ and $t \in]0, 1[$. Define $S_{j,\ell}(y) := \overline{B}_{\mathbb{R}^n}(2^{-j}\ell; 2^{-j}r_1 + y)$. where $j \in \mathbb{Z}$, $\ell \in \mathbb{Z}^n$, and $y \in \mathbb{R}_0$. Suppose that $t_1 \in]0, 1[$. We have

$$\text{supp}_{\text{set}} \Delta_{\mathbf{h}}^m g_k \subset S_{j_0+k, \eta(k)}(m) \quad (7.15)$$

for all $\mathbf{h} \in \overline{B}_{\mathbb{R}^n}(0; t_1)$ and $k \in \mathbb{N}$. Furthermore,

$$S_{j_0+k, \eta(k)}(m) \cap S_{j_0+l, \eta(l)}(m) = \emptyset \quad (7.16)$$

for all $k, l \in \mathbb{N}$ and $k \neq l$. Using Equations (7.15) and (7.16) we obtain

$$\omega_\infty^m(f; t_1) \leq \sup\{\omega_\infty^m(g_k; t_1) : k \in \mathbb{N}\}. \quad (7.17)$$

We have $f \in L^\infty(\mathbb{R}^n)$ and by Equations (7.14) and (7.17) $\omega_\infty^m(f; t_1) \leq c_2 t_1^\sigma$. Thus $f \in B_{\infty, \infty}^\sigma(\mathbb{R}^n)$.

Suppose that

$$f = \sum_{k=0}^{\infty} \mathbf{d}[\iota(k)] \psi_{\iota(k)}$$

for some sequence $\mathbf{d} \in \mathbb{C}^{\Omega(n, j_0)}$ where the series converges in $B_{\infty, \infty}^\sigma(\mathbb{R}^n)$. Define

$$\xi_b := \sum_{k=0}^b \mathbf{d}[\iota(k)] \psi_{\iota(k)}$$

where $b \in \mathbb{N}$. Let $b_1 \in \mathbb{N}$. Define

$$j_1 := \max\{j(\iota(k)) \in \mathbb{Z} : k \in Z_0(b_1)\} + 1.$$

There exists $k_1 \in \mathbb{N}$ so that $\iota(k_1) = \alpha(j_1 - j_0)$. Now $j_1 \neq j(\iota(k))$ for all $k \in Z_0(b_1)$ and consequently $\iota(k_1) \neq \iota(k)$ for all $k \in Z_0(b_1)$. It follows that $k_1 > b_1$. Let $k_2 := j_1 - j_0$. Now

$$\iota(k_1) = (\mathbf{e}_1^{[n]}, j_1, c \cdot 2^{k_2} k_2 \mathbf{e}_1^{[n]}).$$

We have

$$\left| \langle \tilde{\psi}_{\iota(k_1)}, f - \xi_{b_1} \rangle \right| \leq \left\| \left(\langle \tilde{\psi}_{\mathbf{s}, j_1, \mathbf{k}}^{[n]}, f - \xi_{b_1} \rangle \right)_{\mathbf{s} \in J_+(n), \mathbf{k} \in \mathbb{Z}^n} \right\|_\infty$$

and $\left| \langle \tilde{\psi}_{\iota(k_1)}, f - \xi_{b_1} \rangle \right| = 2^{-(j_0 + k_2)\sigma}$. Hence

$$\begin{aligned} 1 &\leq 2^{(j_0 + k_2)\sigma} \left\| \left(\langle \tilde{\psi}_{\mathbf{s}, j_1, \mathbf{k}}^{[n]}, f - \xi_{b_1} \rangle \right)_{\mathbf{s} \in J_+(n), \mathbf{k} \in \mathbb{Z}^n} \right\|_\infty \\ &\leq \left\| \left(2^{j\sigma} \left\| \left(\langle \tilde{\psi}_{\mathbf{s}, j, \mathbf{k}}^{[n]}, f - \xi_{b_1} \rangle \right)_{\mathbf{s} \in J_+(n), \mathbf{k} \in \mathbb{Z}^n} \right\|_\infty \right)_{j=j_0}^\infty \right\|_\infty \\ &\leq \|f - \xi_{b_1}\|_{B_{\infty, \infty}^\sigma(\mathbb{R}^n); j_0}^{(w)}. \end{aligned}$$

Thus $\xi_b \not\rightarrow f$ in Banach space $B_{\infty, \infty}^\sigma(\mathbb{R}^n)$ as $b \rightarrow \infty$.

Theorem 7.3. *Let $n \in \mathbb{Z}_+$, $\sigma \in \mathbb{R}_+$, $q \in [1, \infty[$, and $j_0 \in \mathbb{Z}$. Let $\varphi^{[n]}$ be a mother scaling function of a compactly supported tensor product MRA of $C_0(\mathbb{R}^n)$. Suppose that $\varphi^{[n]} \in C^r(\mathbb{R}^n)$ for some $r \in \mathbb{R}_+$, $r > \sigma$. Then $\{\psi_\alpha : \alpha \in \Omega(n, j_0)\}$ is an unconditional basis of Banach space $B_{\infty, q}^\sigma(\mathbb{R}^n) \cap C_0(\mathbb{R}^n)$ equipped with a norm of $B_{\infty, q}^\sigma(\mathbb{R}^n)$ and the coefficient functional corresponding to basis vector ψ_α is $\tilde{\psi}_\alpha$ for each $\alpha \in \Omega(n, j_0)$.*

Proof. Let

$$\begin{aligned} S_j(m) &:= \{(\mathbf{0}_n, j, \mathbf{k}) : \mathbf{k} \in \mathbb{Z}^n, \|\mathbf{k}\|_2 \leq m\}, \quad j \in \mathbb{Z}, m \in \mathbb{N} \\ D_{\mathbf{s},j}(m) &:= \{(\mathbf{s}, j, \mathbf{k}) : \mathbf{k} \in \mathbb{Z}^n, \|\mathbf{k}\|_2 \leq m\}, \quad \mathbf{s} \in \{0, 1\}^n, j \in \mathbb{Z}, m \in \mathbb{N} \\ D_j(m) &:= \bigcup_{\mathbf{s} \in J_+(n)} D_{\mathbf{s},j}(m), \quad j \in \mathbb{Z}, m \in \mathbb{N}. \end{aligned}$$

Let

$$\begin{aligned} m_1 &:= \min\{m \in \mathbb{N} : \forall \mathbf{k} \in \mathbb{Z}^n : (\|\mathbf{k}\|_\infty > m \implies \forall \mathbf{s} \in \{0, 1\}^n : \tilde{g}_{\mathbf{s},\mathbf{k}}^{[n]} = 0)\} \\ m_2 &:= \lceil m_1 \sqrt{n} \rceil \end{aligned}$$

$$A(k) := S_{j_0}(2^{j_0}k) \cup \bigcup_{l=j_0}^{j_0+k-1} D_l(m_2 + 2^{j_0+k}k), \quad k \in \mathbb{Z}_+.$$

Define $\mathbf{s}(\mathbf{a}) = \mathbf{s}'$, $j(\mathbf{a}) = j'$, and $\mathbf{k}(\mathbf{a}) = \mathbf{k}'$ for all $\mathbf{a} = (\mathbf{s}', j', \mathbf{k}')$, $\mathbf{s}' \in \{0, 1\}^n$, $j' \in \mathbb{Z}$, and $\mathbf{k}' \in \mathbb{Z}^n$. Let $f \in B_{\infty,q}^\sigma(\mathbb{R}^n) \cap C_0(\mathbb{R}^n)$. Let $\eta : \mathbb{N} \rightarrow \Omega(n, j_0)$ be a bijection. Let $\beta_\alpha := \langle \tilde{\psi}_\alpha, f \rangle$ for all $\alpha \in \Omega(n, j_0)$. Define

$$\beta_\alpha^{(m)} := \begin{cases} \langle \tilde{\psi}_\alpha, f \rangle; & \alpha \notin \eta[Z_0(m)] \\ 0; & \text{otherwise} \end{cases}$$

for all $m \in \mathbb{N}$ and $\alpha \in \Omega(n, j_0)$. Define

$$\xi_m := \sum_{\alpha \in \Omega(n, j_0)} \beta_\alpha^{(m)} \psi_\alpha$$

for all $m \in \mathbb{N}$. There exists $c_1 \in \mathbb{R}_+$ so that

$$\left| \langle \tilde{\psi}_\alpha, f \rangle \right| \leq c_1 \|f\|_\infty \quad (7.18)$$

for all $f \in C_0(\mathbb{R}^n)$ and $\alpha \in \Omega(n, j_0)$.

Let $h \in \mathbb{R}_+$. Choose $j_1 \in \mathbb{Z}$, $j_1 > \max\{j_0, 0\}$ so that

$$\left\| \left(2^{j\sigma} \left\| \left(\langle \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]}, f \rangle \right)_{\mathbf{s} \in J_+(n), \mathbf{k} \in \mathbb{Z}^n} \right\|_\infty \right)_{j=j_1}^\infty \right\|_q < \frac{h}{4}. \quad (7.19)$$

Choose $r_1 \in \mathbb{R}_+$ so that

$$\sup\{|f(\mathbf{x})| : \mathbf{x} \in \mathbb{R}^n, \|\mathbf{x}\| \geq r_1\} < \frac{h}{2^{n+2+j_1\sigma}(j_1-j_0)c_1} \quad (7.20)$$

and let $m_3 := \max\{j_1 - j_0, \lceil r_1 \rceil\}$.

Choose $m_4 \in \mathbb{Z}_+$ so that $A(m_3) \subset \eta[Z_0(m_4)]$. Let $l \in \mathbb{N}$, $l > m_4$. If $\mathbf{s}(\eta(l)) = \mathbf{0}_n$ we have

$$\left\| (\beta_{\mathbf{0}_n, j_0, \mathbf{k}}^{(m_4)})_{\mathbf{k} \in \mathbb{Z}^n} \right\|_\infty < \frac{h}{4}. \quad (7.21)$$

If $\mathbf{s}(\eta(l)) \neq \mathbf{0}_n$ we have

$$\left\| \left(2^{j\sigma} \left\| (\beta_{\mathbf{s},j,\mathbf{k}}^{(m_4)})_{\mathbf{s} \in J_+(n), \mathbf{k} \in \mathbb{Z}^n} \right\|_\infty \right)_{j=j_0}^{j_1-1} \right\|_q (j_0 + \mathbb{N}) < \frac{1}{4}h. \quad (7.22)$$

Thus by Equations (7.21) and (7.22) we get

$$\left\| \left(2^{j\sigma} \left\| \left(\beta_{\mathbf{s},j,\mathbf{k}}^{(m_4)} \right)_{\mathbf{s} \in J_+(n), \mathbf{k} \in \mathbb{Z}^n} \right\|_{\infty} \right)_{j=j_0}^{\infty} \right\|_{l^q(j_0 + \mathbb{N})} < \frac{h}{2}. \quad (7.23)$$

By Equations (7.21) and (7.23) we have $\|\xi_{m_4}\|_{B_{\infty,q}^{\sigma}(\mathbb{R}^n);j_0}^{(w)} < h$. It follows that

$$s_m := \sum_{l=0}^m \beta_{\eta(l)} \psi_{\eta(l)} \rightarrow g$$

as $m \rightarrow \infty$ where the series converges in Banach space $B_{\infty,q}^{\sigma}(\mathbb{R}^n)$. We also have $s_m \rightarrow g$ as $m \rightarrow \infty$ in Banach space $C_0(\mathbb{R}^n)$. If we had $g \neq f$ we would have $\langle \tilde{\psi}_{\gamma}, f-g \rangle = \langle \tilde{\psi}_{\gamma}, f \rangle - \langle \tilde{\psi}_{\gamma}, g \rangle \neq 0$ for some $\gamma \in \Omega(n, j_0)$. Now $\langle \tilde{\psi}_{\gamma}, g \rangle = \beta_{\gamma} = \langle \tilde{\psi}_{\gamma}, f \rangle$, which is a contradiction. Hence $f = g$. \square

Theorem 7.4. *Let $n \in \mathbb{Z}_+$ and $j_0 \in \mathbb{Z}$. Let $\sigma \in \mathbb{R}_+$, $p \in [1, \infty]$, and $q \in [1, \infty]$. Let $\varphi^{[n]}$ be a mother scaling function of a compactly supported tensor product MRA of $C_u(\mathbb{R}^n)$. Suppose that $\varphi^{[n]} \in C^r(\mathbb{R}^n)$ for some $r \in \mathbb{R}_+$, $r > \sigma$. Let $f \in B_{p,q}^{\sigma}(\mathbb{R}^n)$. Then*

$$f(\mathbf{x}) = \sum_{\alpha \in \Omega(n, j_0)} \langle \tilde{\psi}_{\alpha}, f \rangle \psi_{\alpha}(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{R}^n$ and the series above converges absolutely for each $\mathbf{x} \in \mathbb{R}^n$.

Proof. Let

$$a := \left\| \left(2^{(\sigma - \frac{n}{p})j} \left\| \left(\langle \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]}, f \rangle \right)_{\mathbf{s} \in J_+(n), \mathbf{k} \in \mathbb{Z}^n} \right\|_p \right)_{j=j_0}^{\infty} \right\|_q.$$

Now

$$\left| \langle \tilde{\psi}_{\mathbf{s},j,\mathbf{k}}^{[n]}, f \rangle \right| \leq 2^{-(\sigma - \frac{n}{p})j} a \quad (7.24)$$

for all $\mathbf{s} \in J_+(n)$, $j \in \mathbb{N} + j_0$, and $\mathbf{k} \in \mathbb{Z}^n$. Let

$$A_0(\mathbf{x}) := \left\{ (\mathbf{0}_n, j_0, \mathbf{k}) : \mathbf{x} \in \text{supp } \varphi_{j_0, \mathbf{k}}^{[n]}, \mathbf{k} \in \mathbb{Z}^n \right\}$$

for each $\mathbf{x} \in \mathbb{R}^n$ and

$$A_k(\mathbf{x}) := \left\{ (\mathbf{s}, j_0 + k - 1, \mathbf{k}) : \mathbf{x} \in \text{supp } \psi_{\mathbf{s}, j_0 + k - 1, \mathbf{k}}^{[n]}, \mathbf{s} \in J_+(n), \mathbf{k} \in \mathbb{Z}^n \right\}$$

for each $\mathbf{x} \in \mathbb{R}^n$ and $k \in \mathbb{Z}_+$. Let $\eta: \mathbb{N} \rightarrow \Omega(n, j_0)$ be a bijection and define

$$\beta_{\alpha}^{(m)} := \begin{cases} \langle \tilde{\psi}_{\alpha}, f \rangle; & \alpha \notin \eta[Z_0(m)] \\ 0; & \alpha \in \eta[Z_0(m)] \end{cases} \quad (7.25)$$

for each $m \in \mathbb{N}$ and $\alpha \in \Omega(n, j_0)$. Let

$$g_m(\mathbf{x}) := \sum_{k=0}^m \langle \tilde{\psi}_{\eta(k)}, f \rangle \psi_{\eta(k)}(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{R}^n$ and $m \in \mathbb{N}$. It follows from Theorem 4.3 that

$$f(\mathbf{x}) = \sum_{k=0}^{\infty} \sum_{\alpha \in A_k(\mathbf{x})} \langle \tilde{\psi}_\alpha, f \rangle \psi_\alpha(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{R}^n$. Define $c_1 := \max\{\|\psi_{\mathbf{s}}^{[m]}\|_\infty : \mathbf{s} \in \{0, 1\}^n\}$. Let $\mathbf{y} \in \mathbb{R}^n$ and $h \in \mathbb{R}_+$. Define $m_1 := \max\{\#A_k(\mathbf{x}) : \mathbf{x} \in \mathbb{R}^n\}$. Let

$$j_1 := \max \left\{ j_0 + 1, \left\lceil \left(\sigma - \frac{n}{p} \right)^{-1} \log_2 \frac{c_1 m_1 a}{1 - 2^{-(\sigma - \frac{n}{p})h}} \right\rceil \right\}$$

Choose $m_2 \in \mathbb{N}$ so that $A_k(\mathbf{y}) \subset \eta[Z_0(m_2)]$ for all $k \in Z_0(j_1 - j_0)$. Suppose that $m \in \mathbb{N}$, $m > m_2$, and $\alpha_0 \in A_{j_0+k}$ for some $k_0 \in Z_0(j_1 - j_0)$. By Equations (7.24) and (7.25) we have

$$\begin{aligned} |f(\mathbf{y}) - g_m(\mathbf{y})| &\leq c_1 \sum_{k=0}^{\infty} \sum_{\alpha \in A_k(\mathbf{y})} |\beta_\alpha^{(m)}| = c_1 \sum_{k=j_1-j_0+1}^{\infty} \sum_{\alpha \in A_k(\mathbf{y})} |\beta_\alpha^{(m)}| \\ &\leq c_1 m_1 a \cdot 2^{-j_1(\sigma - \frac{n}{p})} \frac{1}{1 - 2^{-(\sigma - \frac{n}{p})h}} < h \end{aligned}$$

Thus $g_m(\mathbf{x}) \rightarrow f(\mathbf{x})$ as $m \rightarrow \infty$ for all $\mathbf{x} \in \mathbb{R}^n$. □

References

1. A. Almeida, Wavelet bases in generalized Besov spaces, *J. Math. Anal. Appl.* **304** (2005) 198–211.
2. C. Bennett and R. Sharpley, *Interpolation of Operators* (Academic Press, Inc., London, 1988).
3. P. Brenner, *Besov spaces and applications to difference methods for initial value problems* (Springer, Berlin, 1975).
4. J. Brodzki and G. A. Niblo, Rapid decay and the metric approximation property, arXiv:math.GR/0403423 v2 (2004).
5. C. K. Chui and C. Li, A general framework of multivariate wavelets with duals, *Appl. Comput. Harmon. Anal.* **1** (1994) 368–390.
6. C. K. Chui and C. Li, Dyadic affine decompositions and functional wavelet transforms, *SIAM J. Math. Anal.* **27**(3) (1996) 865–890.
7. S. Dahlke, K. Gröchenig and P. Maass, A new approach to interpolating scaling functions: Scaling functions, *Applicable Analysis* **72**(3-4) (1999) 485–500.
8. I. Daubechies, *Ten Lectures on Wavelets* (SIAM, 1992).
9. I. Daubechies, Orthonormal Bases of Compactly Supported Wavelets, *Commun. Pure Appl. Math.* **XLI** (1988) 909–996.
10. M. Daws, Dual Banach algebras: representations and injectivity, arXiv:math.FA/0604372 v1 (2006).
11. A. Defant and K. Floret, *Tensor norms and operator ideals* (North-Holland, Amsterdam, 1993).
12. G. Deslauriers and S. Dubuc, Symmetric iterative interpolation processes, *Constr. Approx.* **5**(1) (1989) 49–68.
13. R. A. DeVore, B. Jawerth and B. J. Lucier, Image compression through wavelet transform coding, *IEEE Trans. Inform. Theory* **38**(2) (1992) 719–746.

14. R. A. DeVore and V. A. Popov, Interpolation of Besov spaces, *Trans. Amer. Math. Soc.* **305**(1) (1988) 397–414.
15. P. Domański and M. Lindström, Grothendieck spaces and duals of injective tensor products, *Bull. London Math. Soc.* **28** (1996) 617–626.
16. P. Domański, M. Lindström and G. Schlichtermann, Grothendieck operators on tensor products, *Proc. Amer. Math. Soc.* **125**(8) (1997) 2285–2291.
17. D. L. Donoho, Interpolating Wavelet Transforms, tech. report, Department of Statistics, Stanford University (1992).
18. S. Dubuc, Interpolation through an iterative scheme, *J. Math. Anal. Appl.* **114** (1986) 185–204.
19. H. Glöckner, Tensor products in the category of topological vector spaces are not associative, arXiv:math.FA/0307041 v2 (2004).
20. S. Goedecker, *Wavelets and Their Application for the Solution of Partial Differential Equations in Physics* (Presses Polytechniques et Universitaires Romandes, Lausanne, 1998).
21. A. Grothendieck, Produits tensoriels topologiques et espaces nucléaires, *Mem. Amer. Math. Soc.* **16** (1955).
22. B. Han, Symmetry property and construction of wavelets with a general dilatation matrix, *Linear Algebra Appl.* **353** (2002) 207–225.
23. B. Han and R.-Q. Jia, Optimal Interpolatory Subdivision Schemes in Multidimensional Spaces, *SIAM J. Numer. Anal.* **36**(1) (1998) 105–124.
24. W. Härdle, G. Kerkycharian, D. Picard and A. Tsybakov, Wavelets, Approximation, and Statistical Applications, <http://www.quantlet.com/mdstat/scripts/wav/html/>.
25. W. He and M. Lai, Bivariate box spline wavelets in Sobolev spaces, *SPIE's International Symposium on Optical Science, Engineering, and Instrumentation*, (International Society for Optics and Photonics, 1998), pp. 56–66.
26. T. Höynälänmaa, Multiresolution Analysis for Compactly Supported Interpolating Tensor Product Wavelets (long version), arXiv:1005.3371 [math.FA] (2010).
27. H. Ji and Z. Shen, Compactly supported (bi)orthogonal wavelets generated by interpolatory refinable functions, *Adv. Comput. Math.* **11**(1) (1999) 81–104.
28. R. Q. Jia and C. A. Micchelli, Using the refinement equations for the construction of pre-wavelets II: Powers of two, *Curves and surfaces* (1991) 209–246.
29. S. Karakaz'yan, M. Skopina and M. Tchobanou, Symmetric multivariate wavelets, *Int. J. Wavelets Multiresolut. Inf. Process.* **7**(3) (2008) 1–28.
30. J. Kovačević and W. Sweldens, Wavelet families of increasing order in arbitrary dimensions, *IEEE Trans. Image Process.* **9**(3) (2000) 480–496.
31. A. Krivoshein and M. Skopina, Approximation by frame-like wavelet systems, *Appl. Comput. Harmon. Anal.* **31**(3) (2011) 410–428.
32. J. Kustermans and S. Vaes, The operator algebra approach to quantum groups, *Proc. Natl. Acad. Sci. U S A* **97**(2) (2000) p. 547.
33. G. Kyriazis and P. Petrushev, New bases for Triebel-Lizorkin and Besov spaces, *Trans. Amer. Math. Soc.* **354**(2) (2002) 749–776.
34. H. E. Lacey, *The Isometric Theory of Classical Banach Spaces* (Springer-Verlag, 1974).
35. W. Lawton, S. L. Lee and Z. Shen, Stability and orthonormality of multivariate refinable functions, *SIAM J. Math. Anal.* **28**(4) (1997) 999–1014.
36. R. M. Lewis, Cardinal interpolating multiresolutions, *J. Approx. Theory* **76** (1994) 177–202.
37. W. A. Light and E. W. Cheney, *Approximation Theory in Tensor Product Spaces* (Springer-Verlag, 1985).
38. Y. Meyer, *Wavelets and operators* (Cambridge University Press, 1992).

39. P. W. Michor, *Functors and Categories of Banach Spaces* (Springer-Verlag, 1978).
40. O. I. Reinov, On Banach spaces without the approximation property, arXiv:math.FA/0203146 v1 (2002).
41. R. A. Ryan, *Introduction to Tensor Products of Banach Spaces* (Springer, New York, 2002).
42. H. H. Schaefer, *Topological Vector Spaces* (The Macmillan Company, New York, 1966).
43. P.-L. Shui, Z. Bao and X.-D. Zhang, M -band Compactly Supported Orthogonal Symmetric Interpolating Scaling Functions, *IEEE Trans. Signal Process.* **49**(8) (2001) 1704–1713.
44. H. Triebel, *Theory of Function Spaces* (Birkhäuser Verlag, Basel, 1983).
45. P. Wojtaszczyk, *A Mathematical Introduction to Wavelets* (Cambridge University Press, 1997).